

South African Medical Journal

Organ of the Medical Association of South Africa



S.-A. Tydskrif vir Geneeskunde

Vakblad van die Mediese Vereniging van Suid-Afrika

Incorporating the South African Medical Record and the Medical Journal of South Africa

REGISTERED AT THE GENERAL POST OFFICE AS A NEWSPAPER

Vol. 25, No. 38

Cape Town, 22 September 1951

Weekly 2s

IN THIS ISSUE

Van die Redaksie: Editorial

Onhoorbare Klankgolwe en Behandeling

Ultrasonic Waves and Treatment

Original Articles

Education of the Prospective Medical Student

Bilharziasis in the Transvaal

Cranio-Cerebral Trauma

Infantile Eczema

New Preparations and Appliances

Abstracts

Passing Events : In die Verbygaan

In Memoriam

Support your Own Agency Department (P. xviii)

Ondersteun u Eie Agentskap-Afdeling (Bl. xviii)

Professional Appointments (Pp. v, xviii)

for intestinal antiseptis



A highly active sulphonamide of low toxicity for the treatment of gastro-intestinal infections and for prophylaxis during intestinal surgery.



MAY & BAKER LTD

distributors

MAYBAKER (SOUTH AFRICA) (PTY.) LTD., P.O. BOX 1130, PORT ELIZABETH

MA48606

Terramycin

CRYSTALLINE
HYDROCHLORIDE

*Effective
in a wide range
of localized infections...*

1. As the sole therapeutic measure
2. As an adjunct to surgical intervention


Because of its broad antimicrobial spectrum and the promptness of response, Terramycin is now playing an increasingly valuable role in the treatment of localized soft-tissue infections. Erysipelas, erythema multiforme of the bullous type, pyoderma, buccal and penile fusospirochetosis, impetigo, carbuncles, cellulitis, furunculosis and abscesses are among the conditions in which response, described in many instances as "excellent", "good", and "prompt", has been obtained.^{1,2,3,4,5,6} In cases requiring surgical intervention, the use of Terramycin has proved a valuable adjunctive procedure.^{2,3}

Terramycin is available as...


CAPSULES, 250 mg., bottles of 16 and 100; 100 mg., bottles of 25 and 100; 50 mg., bottles of 25 and 100; ELIXIR (formerly Terrabon), 1.5 Gm. with 1 fl. oz. of diluent; ORAL DROPS, 2.0 Gm. with 10 cc. of diluent, and calibrated dropper; INTRAVENOUS, 10 cc. vial, 250 mg.; 20 cc. vial, 500 mg.; OINTMENT, 30 mg. per Gm. ointment; tubes of 1/2 and 1 oz.; OPHTHALMIC OINTMENT, 5 mg. per Gm. ointment; tubes of 1/8 oz.; OPHTHALMIC SOLUTION, 5 cc. dropper-vials, 25 mg. for preparation of topical solutions; TROCHES, 15 mg. each troche; packages of 24.

1. Knight, V.: New York State J. Med. 50:2173 (Sept. 15) 1950.
2. Herrell, W. E.; Heilman, F. R., and Wellman, W. E.: Ann. New York Acad. Sc. 53:446 (Sept. 15) 1950.
3. Blake, F. G.; Friou, G. J., and Wagner, R. R.: Yale J. Biol. and Med. 22:495 (July) 1950.
4. Perdrup, A.: Ugeskrift for Læger 112:1089 (Aug. 3) 1950.
5. Pulaski, E. J.: Ann. New York Acad. Sc. 53:347 (Sept. 15) 1950.
6. Bickel, G., and Plattner, H.: Schweiz. med. Wchnschr. 81:1 (Jan. 6) 1951.

Export Department
CHAS. PFIZER & CO., INC.
81 Maiden Lane, New York 38, N. Y.



Distributors:
Petersen Ltd.
P.O. Box 38
Cape Town, South Africa



How many?

of your patients are in need of
"B" COMPLEX THERAPY

PETERVITE "B" TABLETS

Thiamine HCl 2.0 mgm.
 Riboflavine 1.5 mgm.
 Calc.
 Pantothenate 2.5 mgm.
 Pyridoxine HCl 0.25 mgm.
 Nicotinamide 20.0 mgm.

*In a chocolate-coated tablet
 20's, 60's, 500's*

PETERVITE COMPOUND ELIXIR

Thiamine HCl 1.5 mgm.
 Riboflavine 0.5 mgm.
 Pyridoxine HCl 0.25 mgm.
 Nicotinamide 5.0 mgm.
 Liver Extract (concentrate) 15%

*In TWO teaspoonfuls of
 pleasantly flavoured malt
 base. 4 oz., 16 oz., 80 oz.*

PETERVITE "B" COMPOUND INJECTION

Each 2 c.c. ampoule contains:—
 Aneurine HCl 10 mgm. Riboflavine 2 mgm.
 Nicotinamide 100 mgm. Pyridoxine HCl 5 mgm.
 Calcium Pantothenate 5 mgm.

Box of 6 x 2 c.c.

MADE FOR SOUTH AFRICAN REQUIREMENTS BY



Box 38, Cape Town

Established 1842

Box 5992, Johannesburg

P. 13

BIDUPAN

(formerly Intestinal Concentrated)

FOR RAPID, SUSTAINED, FOUR-WAY RELIEF IN:

- **INTESTINAL INDIGESTION** ● **GALLBLADDER STASIS**
- **BILIOUSNESS** ● **RECURRENT FLATULENCE.**
- etc., etc.

Contains PURE BILE SALTS
 CONCENTRATED PANCREATIN
 DUODENAL SUBSTANCE
 CHARCOAL

IMPROVES BILIARY DRAINAGE and DIGESTION of
 ALBUMIN, CARBOHYDRATES and FATS
 STIMULATES PANCREATIC SECRETION
 REMOVES FERMENTATIVE FACTORS
 SPREADS RELIEF IN BILIOUSNESS, INTESTINAL
 INDIGESTION and RECURRENT FLATULENCE

- SUPPLIED IN BOTTLES OF 50 AND 100 TABLETS; ●
- SUPPLIES AND FURTHER INFORMATION FROM
 OUR DISTRIBUTORS IN SOUTH AFRICA

SIVE BROS. KARNOVSKY LTD.
 JOHANNESBURG and DURBAN

CAVENDISH CHEMICAL CO. (NEW YORK) LTD.,
 OXFORD WORKS, WORSLEY BRIDGE ROAD, LONDON, S.E.26.

ANÆSTHETIC ETHER

Manufactured by
THE NATAL CANE BY-PRODUCTS LTD.
 OF MEREBANK

● Guaranteed to conform to
 the requirements of the 1948
 British Pharmacopœia and the Spec-
 ification of the South African Bureau
 of Standards. Equal to the finest
 imported Ether.

● In cases, each containing
 12 x 1 lb. Amber Coloured Bottles,
 similar to those used in Europe.

For further information please write to the selling Agents

C. G. SMITH & CO. LTD.
 301 Smith Street, P.O. Box 43, Durban

Hert Mendelsohn (Pty.) Ltd.,
 P.O. Box 565, Johannesburg.

C. G. Smith & Co., Ltd.,
 P.O. Box 1314, Cape Town.

Courlanders' Agencies,
 P.O. Box 352, East London.

SECONDARY AMENORRHOEA

Simply and effectively

TREATED ORALLY

WITH

MENSTROGEN

A TYPICAL DOSAGE SCHEME

| | JANUARY | | | | | FEBRUARY | | | | | MARCH | | | | |
|---|---------|---|----|----|----|----------|----|----|----|----|-------|----|----|----|----|
| S | | 3 | 10 | 17 | 24 | | 7 | 14 | 21 | 28 | | 7 | 14 | 21 | 28 |
| M | | 4 | 11 | 18 | 25 | 1 | 8 | 15 | 22 | | 1 | 8 | 15 | 22 | 29 |
| T | | 5 | 12 | 19 | 26 | 2 | 9 | 16 | 23 | | 2 | 9 | 16 | 23 | 30 |
| W | | 6 | 13 | 20 | 27 | 3 | 10 | 17 | 24 | | 3 | 10 | 17 | 24 | 31 |
| T | | 7 | 14 | 21 | 28 | 4 | 11 | 18 | 25 | | 4 | 11 | 18 | 25 | |
| F | 1 | 8 | 15 | 22 | 29 | 5 | 12 | 19 | 26 | | 5 | 12 | 19 | 26 | |
| S | 2 | 9 | 16 | 23 | 30 | 6 | 13 | 20 | 27 | | 6 | 13 | 20 | 27 | |

The dosage is 4 tablets daily for 5 days.

Menstruation under this treatment may be expected to follow in 4 to 6 days. Certain cases of habitual abortion will also respond to this treatment.

Concise but full information gladly sent on request.

○—4 Tablets Daily — For 5 Days — Every 4 Weeks

□—Menstruation can be expected

MENSTROGEN

Each tablet contains:

LYNORAL (ETHINYL (ESTRADIOL) 0.01 mg.

PROGESTORAL (ETHISTERONE) 10 mg.

In tubes of 20. Bottles of 60 tablets.



ORGANON LABORATORIES LTD., LONDON

Literature on request from

SOLE SOUTH AFRICAN DISTRIBUTORS

KEATING'S PHARMACY LTD.

P.O. Box 256, JOHANNESBURG

P.O. Box 568,
CAPE TOWN

P.O. Box 2383,
DURBAN

P.O. Box 931,
PORT ELIZABETH

South African Medical Journal Suid-Afrikaanse Tydskrif vir Geneeskunde

Vol. 25, No. 38

Cape Town, 22 September 1951

Weekly 2s

CONTENTS

| | | | |
|--|-----|--|-----|
| The Education of the Prospective Medical Student. Mr. T. Lindsay Sandes, F.R.C.S. | 685 | Bilharziasis in the Transvaal (Concluded from p. 680). Dr. S. Annecke and Dr. P. N. B. Peacock | 689 |
| New Preparations and Appliances: Tenamid—A New Anthelmintic | 687 | Aspects of Cranio-Cerebral Trauma: I. The Surgical Anatomy and Physiology of Cranio-Cerebral Trauma. Dr. D. A. Muskat | 692 |
| Abstracts | 687 | Infantile Eczema. Dr. S. Gordon | 697 |
| Van die Redaksie: Onhaarbare Klankgolwe en Behandeling | 688 | In Memoriam: Dr. J. S. Morton | 700 |
| Editorial: Ultrasonic Waves and Treatment | 688 | Passing Events: In die Verbygaan | 700 |

Leocillin

THE NEW ANTIBIOTIC

Benzylpenicillin-diethylaminoethyl ester hydriodide

The ANTIBIOTIC

for

*selective penicillin therapy
in lung tissues*



PHARMAKERS (PTY.) LTD.

215-6 Gibraltar House, Regent Road, Sea Point, CAPE TOWN

Registered Agents for

LEO PHARMACEUTICAL PRODUCTS COPENHAGEN

(5)

Partnership Required

Young married doctor, fully bilingual, with hospital and general practitioner experience, seeks partnership in Cape Town or any town in Western Cape Province. Write to 'A. I. D.', P.O. Box 643, Cape Town.

Required

Medical officer required for Gath's Mine (Asbestos), Mashaba, Southern Rhodesia, immediately. Salary £1,200 per annum, with free house and transport provided. Further particulars on application to Principal Medical Officer, Shabanie Mine, Shabani, Southern Rhodesia.

For Sale

One Klinostik Diagnostic set and one doctor's bag. For further information write to the Local Agency Manager, Medical Association of South Africa, 112 Medical Centre, Field Street, Durban.

DOCTORS' BAGS in GENUINE BRITISH LEATHER


Brown Crocodile finish

Two locks and keys

Leather Handle

Five drawers

Space for Sphyg.



As illustrated,
but extra drawer instead of Urine Set

UNBEATABLE VALUE

at £12-10-0

Ask for Model 'Webley'

at



Union Medical Supplies

81 Long Street, Box 3907 Tel. 2-2955
CAPE TOWN

E.N.T. Outfit

Retiring E.N.T. Surgeon offers at bargain price complete set of E.N.T. instruments and apparatus, including electric ear masser, electric pharyngoscope, Haslinger bronchoscope and oesophagoscope (Vienna), audiometer (Pillings, Philadelphia), bougies, dilators, etc. Five-shelf white enamelled instrument cabinet, etc. Original cost over £600. Bargain at £300. Apply 'E.N.T.', P.O. Box 209, Maritzburg, Natal.


Assistantship Required

An assistantship is required with a general practitioner in Cape Town, Muizenberg or vicinity. Applicant qualified in 1944. Write to P.O. Box 4627, Cape Town.

Johannesburg Partnership Wanted

Experienced Jewish general practitioner desiring to move from large country practice to Johannesburg seeks partnership in good class practice. Write to 'A. I. E.', P.O. Box 643, Cape Town.

'Sulphamezathine' in Pneumonia



"These observations confirm that 'sulphamezathine' is highly satisfactory in lobar pneumonia . . . Its lower toxicity may help to reduce the mortality in very ill patients, while the virtual absence of vomiting ensures more complete absorption."—*Lancet*, 1944, i, 277.

"Of the commonly available sulphonamides, 'Sulphamezathine' causes relatively little nausea and vomiting and, having an acetyl derivative of high solubility, very rarely gives rise to renal complications. It is therefore suitable for routine treatment."—*British Medical Journal*, 1949, ii, 1225.

IN CHILDREN: "The results of treatment with 'Sulphamezathine' therefore compare favourably . . . showing the lowest mortality rate, the most rapid response to treatment and freedom from toxic effects."—*Arch. Dis. Childh.*, 1944, 19, 122.



'SULPHAMEZATHINE'

SULPHADIMIDINE B.P.C.

TRADE MARK

the safe sulphonamide

A product of

IMPERIAL CHEMICAL (PHARMACEUTICALS) LIMITED

(A subsidiary company of Imperial Chemical Industries Ltd.)

WILMSLOW, MANCHESTER

Distributed by

I.C.I. South Africa (Pharmaceuticals) Limited · P.O. Box 7796, Johannesburg

South African Medical Journal

Suid-Afrikaanse Tydskrif vir Geneeskunde

Vol. 25 No. 38

Cape Town, 22 September 1951

Weekly 2s

THE EDUCATION OF THE PROSPECTIVE MEDICAL STUDENT*

T. LINDSAY SANDES, F.R.C.S.

Cape Town

Education might be defined as a knowledge of facts, and of methods and conduct, in preparation for the work of living. The major part of education was directed towards providing a living or bringing in an income. One might call this utilitarian education. On the other hand, the work of living did not cease when one shut one's shop, or office, or upon laying down the stethoscope or ophthalmoscope. One owed a duty to one's family, to the community and to the State. Education should make some preparation for these responsibilities. Owing to the length, the difficulty and the intensity of medical education, in a relative assessment of these two departments of education, an increasing want of equilibrium or balance was becoming apparent. Our medical schools were undoubtedly now, on the whole, turning out well-trained young students as doctors. This was excellent as far as it went, but it was having a serious reciprocal effect. There is a fear that the standard of general education was now becoming deficient. This deficiency will in time affect our prestige as a profession and our authority as doctors. Now consider for a moment the upbringing of a boy at school. In his early years he receives a good basic preliminary education; in his later years, certainly in many schools of the Old Country, there has been introduced of recent years a pernicious system of dividing the boys in the senior classes into two categories—the classical side and the modern side. On the classical side are collected the boys who are, for example, to become theological students intended for law, journalism, or for posts in the higher branches of the civil service. These boys study for the most part literary subjects, such as history, philosophy, the ancient classics. On the other hand, the boys destined to become chemists, engineers, geologists and doctors—their studies are mainly directed to science and mathematics. In consequence, the proper basis of education, a reasonably comprehensive foundation of knowledge, has been infringed; at an early age these children are directed towards becoming premature specialists with the narrow, the constricted, the prejudiced outlook that specializing tends to engender. The boy now passes his matriculation examination. This is a minimum standard of education for admission to the

University. It is a standard which an average typist attains, a clerk in a Government office, a bank, or an insurance office, and many probationer nurses; yet it seems to be all that is required of students for the learned profession of medicine. He now enters the University but instead of mixing and communing with other undergraduates, instead of imbibing the University atmosphere, instead of getting the advantages of a broad and mixed undergraduate society of students of all types of all the professional sections and of the Arts, he at once enters the Medical School and lives in an isolated medical residence. From thence onwards he is immured in a medical monastery.

Here for the next six or seven years he perambulates the cloistered walks of the Institution. He hears nothing, he sees nothing, he inhales nothing but medical topics of one sort or another. He forgets there is a world outside. He is hardly cognizant of its existence; during those six or seven years the fleshy tablets of his mind are receiving nothing but a torrent of medical impressions. Those fleshy tablets, wax to receive and marble to retain, are permanently recording that which will affect and direct his views, his associations, and his conduct of affairs for the rest of his life. He will observe life through a polarized medium; he will see with a mental astigmatism. If he proceeds to specialize in a particular department of medicine, this imbalance of his education becomes one of a still more unstable equilibrium.

What methods can be recommended to help to correct this want of balance in his educational diet? Much as one dislikes to suggest it, after admission to the University, the prospective medical student should do one year of basic general education; in that year the subjects he studies should have no relation to, or be in any sense ancillary to, those of his medical curriculum. One recommends particularly the humanities such as history, literature, philosophy, logic and languages. I have a particular prejudice in favour of the ancient classics. This year in addition would allow a better selection of students for admission to the medical school and those who obviously had no vocation for such a professional life could more easily be separated. Furthermore, judging from 25 years' experience as a teacher of medical students, I feel less reluctant in recommending this preliminary year because those students who had a better basic education, or who were a year or two older than the average of the

* Report of the Presidential Address delivered at a meeting of the Cape Town Post-Graduate Medical Association on Thursday evening, 19 April 1951.

class, or who before had had some educational experience in other lines such as teachers, seem to me to have been better students, did better at their examinations, and made a greater success of their professional life in later years. In fact, such a year or preliminary educational years were not lost. They constituted a credit reservoir of learning and experience that was of immense advantage in the preparation for life and in practice as a doctor. I admit that this might constitute an economic hardship but I would be prepared to accept for this preliminary education a year in any employment of an educational nature which might even bring in an emolument, e.g. in place of that year, a five-months' stay in the Antarctic on a whaler or, if it could be arranged, the student should go through a course at Voortrekkerhoogte in training as a policeman, or a year in the Defence Force, or a year's experience in a foreign country where he learnt the language, or, indeed, in any employment, salaried or otherwise, provided it could be demonstrated that it was some type of educational employment, or where he enlarged his experience of men and affairs or as a citizen of the world.

The biographies of some great and successful men demonstrate how such preliminary experience not only helped but was almost a major factor in making a success of certain careers. Sir Charles James Napier, whose statue adorns Trafalgar Square, at the age of 14, was bought a commission as an ensign in the Army; his colonel, contrary to custom but being a friend of his father, sent him back home to Ireland for another four years' general education before being trained as an officer in the Army; he was severely wounded in the Peninsular wars where he lost an eye. Napier became a great general, a first-class soldier, he served with great distinction in Baluchistan, and was later a good administrator. He maintained that the success he made of his career was particularly due to the preliminary non-military education before he got submerged in his duties as a soldier. He was the only General recorded in history (since Julius Caesar) who could make a good pun in Latin.

The life of Lord Reading, the son of a Mr. Isaacs, a penurious fruit broker, is also to the point. Rufus Isaacs, finding that the living and the prospects in his father's fruit broker's office were by no means bright, went off to sea as a sailor before the mast in a windjammer. He sailed the ocean and he braved the elements as a deck-hand in this limejuicer; he became a tough, hardy, weather-beaten seaman. After many adventures the ship dropped anchor in Bombay and Rufus the young sailor got leave ashore while the ship awaited instructions. The instructions to the ship were to return home, and presently this bronzed sailor appeared in his father's office. But things were no better in the office after his years at sea; he told his father that he wanted now to study law. The old man, dismayed and surprised, explained that Rufus had neither the education, nor himself the money to permit him to study law. Rufus replied that he had made full inquiries, he had made some preparations, and he proposed to sit for the examination for the Middle Temple; although he had not much money he had sufficient saved to carry him through part of the legal training. He passed the entrance examination to the Middle Temple. He sat his lectures, he ate his dinners, and at the late age of 27 he was called to the Bar. His progress

was rapid and at the age of 38, 11 years after his admission to the Bar, he took silk—almost a record. Later he entered Parliament, became a Judge, then Lord Chancellor. In due course his sovereign appointed him to the high office of Viceroy of India. Forty years after that sailor boy had climbed the seaweed covered and slippery stone steps of Bombay pier, the gateway of India, he again ascended those steps with all the emblazonry, ceremony and dignity as Viceroy of a great nation of 350 million people. Lord Reading attributed a material proportion of his success to his preliminary experiences, to the hard education he had received while a sailor on the high seas. It gave him a knowledge of men and affairs, to push aside the meretricious and to accept facts. These instances demonstrate that reasonably limited preliminary experience, as long as it was of an educational type, although it might delay the entrance to one's profession, was generally repaid with compound interest in the course of the practice of one's profession. But for the average young medical student straight from school, one year's preliminary literary education was almost a *sine qua non*.

I myself have a preference for certain subjects in this preliminary study of the prospective medical student. I would emphasize my personal regard for the value, as an educational discipline, of the ancient classics, Greek and Latin. These subjects did not seem so frequently taught in these days as they were in past generations. They were often mislabeled dead languages. I would like, for a moment, to recall a little of what we owe to these ancient civilizations.

Greece, in the fifth century B.C., set standards of art, architecture, oratory, philosophy, drama, literature, logics, historical records, science and scientific research, poetry, and principles of democracy that have rarely been approached and never exceeded since those long ago times. Ancient Greece gave us the nucleus and the substance of the western civilization on which the best of modern scientific and ethical progress depends. When Greece was overwhelmed by the growing Roman Empire, that stern military administration recognized the great essential value of Grecian civilization. Rome emulated and absorbed it in its own life and institutions. Furthermore, the sons of the nobles of the Roman Empire and of those who could afford it, were sent to Greece to learn the language and to absorb the culture which the Romans deeply appreciated. So that through the Roman Empire we have received and impounded a great deal of the culture and the civilization that was Grecian and on which the West is founded. The Roman Empire lasted 1,000 years. It extended from the lowlands of Scotland in the west to the Tigris and Euphrates in the east; from the Danube, the Rhine and the Baltic in the north. It embraced the North African littoral to the burning deserts of the south. The engineering of Rome, its buildings, bridges, its roads and highways, are there to be admired to this day. Roads that travel almost in straight lines from the capital to the remotest marches of the Empire; over mountains, across valleys, by bridges and viaducts. Communication was swift and efficient. Roman law first enrolled on the 12 tables in the early days of the Roman kings and later enlarged and amended by the Republic. Ultimately Justinian in A.D. 529, after 1,300 years of legislation,

promulgated an imperial statute comprehending a simplified code and digest of valid laws. Those laws form the basis of the legal systems of all Southern Europe, the British Isles and of all the Americas. Again in the fortified outposts of the Empire the Roman soldiers were good ambassadors and agents. The people and the peasantry, whether in Spain or Portugal or Gaul or Belgium or Italy, learnt to speak the soldiers' and traders' Latin. It was not the classical Latin of Cicero's orations, nor the sonorous phrases of Virgil's hexameters. It was the common speech of the ordinary people. In each of these countries this speech was gradually adopted, gradually spread and adapted, affected and modified by local intonation and dialect and developed into the languages of each of these nations. Half the words of English which I myself am now using were derived directly or indirectly from Latin; and the civilized speech of all the Americas, from Canada to Magellan, was either a modified Latin through the Spanish or Portuguese, or in the English-speaking parts, half is derived from Latin. The civilization of ancient Rome in which the Grecian standards were absorbed was at a very high level in the best days of the Empire. It was said that no Roman citizen was illiterate. He could read and write and figure. He had a definite standard of education; the only members of the Roman Empire who did not rise to those standards were the lowest class of slaves. The better class of slaves were often highly educated and acted as teachers and tutors. But the lowest class of slaves, e.g. coolies and labourers, did not have this standard nor did the recently conquered barbarians. The very word barbarian is imitative of the rude speech of a wild, illiterate race. It was a highly prized and privileged honour, but not without its obligations, to be able to say *Civis Romanus sum*.

About the fifth century A.D. the furious and repeated inroads of the vandals, the Goths and the Huns commenced to press back the Roman Empire, already enfeebled from many intrinsic ailments. Western civilization commenced to collapse. The Dark Ages had begun.

For four or five centuries it looked as if civilization would be entirely submerged and destroyed; the world, indeed, nearly slipped back to prehomeric standards. But all through those dark ages civilization nearly asphyxiated, buried and hidden, still maintained a spark of life; credit for the nurture and maintenance of that spark of life must be given in the first place to the monasteries—the monasteries of the great Catholic brotherhoods. Their records were kept in classical Latin, their communications and commerce were conducted in that speech and while the Latin language survived it could always bring back the immense documentary records and literature of the lost civilization. About the tenth to the twelfth centuries many of these monasteries began to develop into educational institutions and these were the beginnings of Universities. Padua, Salamanca, Prague, Heidelberg, Paris, Vienna, Oxford, Cambridge, St. Andrews—Salerno and Bologna Universities are said to have the distinction of being the first. The renaissance had dawned. Since then our civilization has developed rapidly. It is founded upon the principles, the precepts, the methods, and the culture derived in the first place from ancient Greece and Rome. The very language of scientists, of Newton, of Harvey, of Gillileo, up to mid-Victorian times, was classical Latin.

I must refer to the pleasure and refreshment that I derive from the well-thumbed textbooks of the classics which I still have from my far-off youth. I admit I am prejudiced in favour of the classics. I am convinced of their basic educational value. Though not so readily convertible into commercial barter as technical scientific attainments, they had an incontrovertible claim to be considered in any educational curriculum. It was for these reasons that I recommend them in any preliminary course before students enter their medical school in an attempt to balance the overweighted scientific studies which the medical student must later undertake.

Such a pre-medical educational year should be founded on the humanities. What shall it profit a man if he gain the whole world and lose his own soul.

NEW PREPARATIONS AND APPLIANCES

TENAMID: A NEW ANTHELMINTIC

Tenamid, the recently discovered anthelmintic preparation of the Schering Corporation, allows for the rapid and effective oral therapy of tenia infestation of the intestines.

Chemically known as β -phenyl- β -carboxy- (3, 5-diiodo-4-hydroxy phenyl) ethane, *Tenamid* has been proved effective against *Taenia solium*, *Taenia nana*, *Taenia saginata*, *Necator americanus*, *Hymenolepis*, *Dipylidium caninum*, *Bothriocephalus latius*, *Trichocephalus* and *Ascaris*, in a high percentage of cases reported.

Usually one course of treatment is sufficient to expel the parasites completely. A second or even a third course of treat-

ment may be instituted at a 6- to 8-day interval, in that rare case which does not respond to a single course.

Tenamid is in tablet form and administration is simple. No special diet or purgatives are necessary before treatment. The successful therapeutic results obtained are marked by a complete absence of detrimental secondary effects.

Tenamid tablets each 0.5 gram are available in tubes of 12 and bottles of 100.

Manufactured in the Union of South Africa by Schering (Pty.) Limited, Johannesburg, for and under the formula and technical supervision of Schering Corporation, Bloomfield, N.J.

ABSTRACTS

Anti-Perniciious Anaemia Factor and White-cell Count. J. Dedichen and P. Laland (1949): *Lancet*, **257**, 282.

As already found with folic acid, pure crystalline Vitamin B_{12} (40 micrograms) had no effect within one day on the leucocyte count in one normal patient. Following a subsequent dose of 10 cc. liver extract, leucocytosis occurred promptly.

Vitamin B₁₂. D. M. Dunlop and W. M. Wilson (1949): *Lancet*, **256**, 754.

An extract from a strain of *Streptomyces* after assaying microbiologically for Vitamin B_{12} activity was used to treat two cases of Addisonian pernicious anaemia. The response was equivalent to that expected from a potent liver extract.

South African Medical Journal

Suid-Afrikaanse Tydskrif vir Geneeskunde

VAN DIE REDAKSIE

ONHOORBARE KLANKGOLWE EN BEHANDELING

Die verwarmende invloed van onhoorbare klankgolwe is biologies gesproke van weinig belang. Van groter belang egter is die eienskap wat dit besit om 'n vermeerderde bloedsrykheid te veroorsaak, terwyl daar ook aanspraak op gemaak word dat hierdie golwe 'n byna spesifieke meganiese invloed besit op die vog in die weefselruimtes, die verspreiding daarvan, met deurgang deur weefselvliese, asook op die gemiese toestande en dié van die struktuur van die sel. Op grond van hierdie eienskappe is daar al aanspraak op gemaak dat gunstige geneeskundige uitlae behaal is by die behandeling van gewrigs-, senuwee-, en veselweefseltoestande. Dit het in die berigte nie ontbreek aan verslae nie, omtrent die goeie subjektiewe gevolge by gevalle van 'ulcus duodeni, vergroeiings in die die buikholte, galblaasontsteking . . . en senuweepyn van die buikwand'.¹ Die invloed van behandeling met onhoorbare klankgolwe is al toegeskryf geword aan mikromassering van die selle sowel as aan verhoging in die temperatuur van die weefsels sowel as aan kolloidaal-veranderinge.

Terwyl daar weinig twyfel kan bestaan dat, onder proef-ondervindelijke omstandighede, hierdie golwe 'n mees diepgaande invloed uitoefen op die weefsels, is daar 'n neiging by die meerderheid van hierdie verslae, waarin daar aanspraak gemaak word op gunstige geneeskundige resultate, om van uit Westelike Europa te kom. In Engelse en Amerikaanse hande was die uitslae nie van bepaalde aard of enig nie en het dus nie aangetoon of hierdie vorm van behandeling (wat geensins sonder gevaar is nie) bepaalde vooruitgang van enige soort verteenwoordig, bo enige van die bestaande wyses van behandeling nie.

DIE GEVAAR

Die onhoorbare golwe mag, wanneer dit gerig word op die plexus brachialis, wederkerende hartkloppings en hartkramp veroorsaak. Alhoewel daar al aanspraak op gemaak is dat goeie uitslae verkry word by die behandeling van sere aan die been, bly daar nóg die gedurige gevaar dat plaaslike bloedklonte in beweging gebring kan word. Die golwe kan ook, as gevolg van hul invloed op die geslagsklier, onvrugbaarheid veroorsaak; hulle kan die ontwikkeling van die groeiende gebeente by kinders hinder en maak dit vir die werker iets gewaagds. Dit is daárom duidelik dat die werktuie vir die vervaardiging van hierdie golwe as uiters gevaarlik beskou moet word. Daar kan geen verskoning voor wees dat leke toegelaat sal word om hierdie dinge te hanteer nie en dit is wenslik dat selfs lede van die mediese beroep die toestel alleenlik onder leensie sal gebruik.

1. Excerpta Medica (1951): Afdeling VI (Internal Medicine), 5, 317.

EDITORIAL

ULTRASONIC WAVES AND TREATMENT

The thermogenic effect of ultrasonic waves is biologically not of much significance. More important is the property of inducing hyperaemia and it has also been claimed that these waves have almost a specific mechanical effect upon the circulation of fluid in the tissue spaces, diffusion through tissue membranes and the chemical and structural conditions in the cells. It is on the basis of these properties that favourable therapeutic results have been claimed in the treatment of arthritic, neuritic and fibrositic conditions. Reports have not been lacking about good subjective results in cases of 'duodenal ulcer, abdominal adhesions, cholecystitis . . . and neuralgia of the abdominal wall'.¹ The effect of ultrasonic therapy has been ascribed to micromassage of the cells as well as an increase in the temperature of the tissues and to colloidal changes.

While there can be little doubt that under experimental conditions these waves exercise a most profound effect on tissues, most of the reports claiming favourable therapeutic results tend to come from Western Europe. In English and American hands the results have not been specific or unique and have not indicated that this form of treatment (which is not without its dangers) represents any advance of any kind on any of the existing methods in current use.

DANGERS

The ultrasonic waves may, when directed to the brachial plexus, produce paroxysmal tachycardia and angina pectoris. Although it has been claimed to have good results in the treatment of ulcers of the leg, there is the ever-present danger of mobilizing regional thrombi. The waves can also, by their action on the gonads, produce sterility; they can interfere with the development of growing bones in children and place the operator of the instrument at risk. It is clear, therefore, that instruments for producing these waves must be regarded as highly dangerous. There can be no excuse for permitting their use by lay persons and it is desirable that even members of the medical profession should use the apparatus only under licence.

1. Excerpta Medica, 1951, Section VI (Internal Medicine), 5, 317.

"TAMPOVAGAN"

(Regd.)

PESSARIES

"Tampovagan" Pessaries are issued as follows:

- Lactic Acid 5%¹⁰
- Ichthylol 5%¹⁰
- Ichthylol 10%¹⁰
- Choleval 1%¹⁰ (Silver Proteinates)
- Stilboestrol and Sulphathiazole
- Penicillin (5,000 i.u.) and
- P.S.S. (Penicillin-Sulphanilamide-Sulphathiazole)

ALL TAMPOVAGAN GLOBULES (except Tampovagan with P.S.S. and with pure penicillin) are of a glycerine-gelatin base, which is non-irritating, dissolves speedily at body temperature and absorbs the oedematous fluid from the tissues, thus reducing the inflammatory state and the rate of discharge, and enhancing the action of the main ingredients.

Tampovagan globules containing penicillin have a cocoa-butter base because penicillin was found to be incompatible with gelatine.

DOSAGE:

One Tampovagan globule to be inserted into the vagina every night before retiring to bed, and in the morning approximately 30 minutes before rising.

PACKINGS: BOXES OF 8

CAMDEN CHEMICAL CO. LTD.

61 Gray's Inn Road, London, W.C.1, England
Telephone: Holborn 7524 Telegrams: "Camkemco, Holb.", London

Sole South African Agents:

WESTDENE PRODUCTS (PTY.) LTD.

22-24 ESSANBY HOUSE, 175 JEPPE STREET, JOHANNESBURG P.O. BOX 7710 PHONE: 23-0314

CAPE TOWN: 408/9 CTC Buildings, Plein Street Phone 2-2276 DURBAN: Alliance Buildings, Gardiners Phone 2-4975
PRETORIA: Central House, Central Street Phone 3-3487 PORT ELIZABETH: P.O. Box 607

(a) **TAMPOVAGAN WITH LACTIC ACID 5%¹⁰**
Indications:
Leucorrhoea due to neglected pessaries, cervical tears, hormonal disturbances, mild infections.

(b) and (c) **TAMPOVAGAN WITH ICHTHYOL 5%¹⁰ AND 10%¹⁰**
Indications:
All degrees of inflammatory changes of the vagina and of the cervix in particular. Tampovagan with ichthylol 10% should be used in strong inflammatory conditions, with ichthylol 5% in milder cases only.

(d) **TAMPOVAGAN WITH CHOLEVAL 1%¹⁰ (SILVER PROTEINATES)**
Indications:
Mild inflammatory changes of the vagina with medium discharge due to mild infection, or as a continuation of therapy after a course of Tampovagan with P.S.S.

(e) **TAMPOVAGAN WITH PENICILLIN (5,000 i.u.)**
Each globule contains 5,000 i.u. of penicillin in a cocoa-butter base.
Indications:
All cases of vaginal fluor in which the causative organism is known to be sensitive to penicillin, in cases showing an idiosyncrasy towards sulpha drugs, or where sulphonamides have been previously employed and have not achieved the desired therapeutic results.

(f) **TAMPOVAGAN P.S.S. (PENICILLIN, SULPHANILAMIDE AND SULPHATHIAZOLE)**
These contain 5,000 i.u. of penicillin with 0.25 g. of sulphanilamide and sulphathiazole.
Indications:
All primary and secondary infections due to organisms sensitive to penicillin and/or sulphonamides. Gonorrhoea, syphilis, streptococcal and pneumococci, etc.

(g) **TAMPOVAGAN WITH STILBOESTROL AND SULPHATHIAZOLE**
Indications:
Semile vaginitis, B. coli infections, hypo-oestrogenised function, etc.



SEDATION IN DYSMENORRHOEA

Dysmenorrhoea is a symptom or entity in which Gelonida* provides prompt and effective relief not only of pain but also of the associated mental distress. The anxiety and irritability so characteristic of genital disturbances is particularly evident in dysmenorrhoea.

Gelonida is both analgesic and sedative and may be confidently prescribed in the treatment of pain and anxiety in menstrual distress.

GELONIDA

Supplied in tubes of 10 and 20 tablets,
also bottles of 50 and 100 tablets.

Distributors: CHAMBERLAIN'S (PTY) LTD., 6-10 Searle Street, Capetown.
Successors to: William R. Warner & Co. Ltd., Power Road, London.

144 Ex

ANTABUS

the new medical treatment for
ALCOHOLISM

IMPORTANT CONSIDERATIONS IN "ANTABUS" TREATMENT.

1. "ANTABUS" is not a cure for Alcoholism, it is an aversion treatment, and its use must be accompanied by careful observation and measures aimed at correction of underlying personality disorders. For this reason, it is essential to obtain the consent of the patient, and where possible, the co-operation of relatives. Follow-up visits and encouragement are of great importance.
2. As in the case of all new therapies, great care must be exercised in patients suffering from Cardiovascular diseases; patients having less than 85% of normal Liver function; chronic or acute Nephritis; Epilepsy; Diabetes mellitus; Asthma and Pregnancy.
3. "ANTABUS" should not be administered to patients who have been given Paraldehyde as it may be metabolised through an Acetaldehyde stage. Similarly Paraldehyde should not be administered to "ANTABUS"-treated patients.
4. The patients desire to stop treatment should be discouraged until such time as it is confidently felt that social readjustment has been effected. The aid of social workers such as "Alcoholics Anonymous" is, in many cases, of great importance.
5. "ANTABUS" is a relatively safe drug provided a proper physical, psychiatric and social evaluation of the patient is made before treatment is commenced.
6. In cases of violent reaction Nikethamide and a mixture of 95% Oxygen and 5% Carbon Dioxide have been administered with good effects.

TRADE ENQUIRIES:

NATAL: Stuart Jones and
David Anderson, Ltd., 20 Queen
Street, Durban.

TRANSVAAL and O.F.S.: B.
Owen Jones, Ltd., 83 Main
Street, Johannesburg.

CAPE, Eastern Province: B.
Owen Jones Ltd., 63 Cambridge
Street, East London.

CAPE, Western Province: Sciex
(B. Owen Jones), Ltd., Raphael's
Buildings, 86 Darling Street,
Cape Town.

Terwyl dit twyfelbaar lyk of daar enige geneeskundige deug bestaan vir die aanwendig van onhoorbare golwe en omdat daar nog geen vaste wetenskaplike saak voor uitgemaak is vir toeskrywing aan hierdie golwe van enige geneeskundige voortreffelijkheid op spesifiekheid nie, behoort ons uiters versigtig te wees t.o.v. die binnelating van hierdie gevaarlike instrumente na die Unie en, veral, die hantering daarvan deur leke. Van welke aard die aansprake ook al sal mag wees, moet hul op hierdie stadium gewend word na die gebied van die navorsing liewers as na die van die kliniese praktyk.

As it seems doubtful whether any special therapeutic virtue is inherent in the use of ultrasonic waves and because no sound scientific case has yet been made out for giving these waves any therapeutic pre-eminence or specificity, we should be most circumspect about the introduction of these dangerous instruments into the Union and, in particular, their use by inexperienced lay persons. Whatever claims are to be made should at this stage still be directed to the field of research rather than clinical practice.

BILHARZIASIS IN THE TRANSVAAL

S. ANNECKE, M.A. (CAPE TOWN), M.B., B.S. (DURHAM), B.Hy. (DURHAM), D.P.H. (LOND.), D.T.M. & H. (LOND.)
M.R.C.S., L.R.C.P.

and

P. N. B. PEACOCK, M.B., Ch.B. (CAPE TOWN), D.P.H. (CAPE TOWN)
Union Health Department, Tzaneen, Transvaal

(Concluded from page 680)

(c) *Distribution of Bilharziasis (S. Haematobium only).* In 1937-1938 a somewhat limited survey to determine the infectivity rate amongst Native school-children was undertaken. The results of this survey appears in Table E.

A much more extensive survey which covered nearly the whole of Transvaal where bilharziasis occurs, followed in 1948. Whereas the 1937-1938 survey dealt primarily with Natives the later one included numbers of European school-children as well.

Attached Tables E and F reflect the distribution of bilharziasis amongst Native school-children in the Transvaal in the years 1937-1938 and 1948-1950 respectively. Table G conveys the same information for European school-children in the years 1948-1950. To simplify these tables a number of schools from each district have been lumped together with the result that the very high percentage incidence (90%+) found in certain individual schools is not shown. (In the 1937-1939 figures, when smaller numbers were examined the higher percentages are more obvious.) The attached map shows those parts of each area where the percentage incidence was above or below the pooled figures given in the latter. Our criterion for considering a subject as having Bilharzias was the microscopical identification of viable Schistosoma ova in a single centrifuged terminal urine specimen collected under supervision after moderate exercise as recommended by Bennie.¹⁷

We notice from the tables that:

i. Bilharzias is widely distributed amongst Native school-children over the Northern and Eastern Transvaal, with infectivity rates of up to 80%.

ii. The surveys of 1948-1950 correspond closely with the 1937-1938 surveys.

iii. The highest incidence of bilharziasis exists in the North and North-Eastern Transvaal Lowveld and portion of the Western areas of the North Transvaal with less in the intervening Highveld (Map, Fig. 1). This map also shows how patchy the distribution of bilharziasis is in the Northern Transvaal.

iv. The incidence of Bilharzias amongst European school-children is much lower than amongst Native school-children, but the occurrence of the disease is as widespread.

It should be noted that those figures only refer to a selected group (i.e. children) and they cannot be considered as giving the general incidence amongst the whole population.

The accompanying map, referred to above, deals only with the distribution of *S. Haematobium* in the Transvaal and shows many areas where more than 60% of the Native school-children are infected (sometimes 60% and over). The distribution of *S. Mansoni* has not been explored to any great extent but it is known to occur fairly frequently with the majority of the known cases concentrated in the Transvaal Lowveld to the east of the Drakensberg Mountains.

TABLE E: URINE EXAMINATIONS NATIVE SCHOOL-GOING CHILDREN 1937-1938

| Zone | Locality | Number Examined | % Positive |
|---------|--------------------|-----------------|------------|
| Zone I | Shulivane | 322 | 67 |
| | Sibasa | 1,577 | 31 |
| Zone IV | Nelspruit | 119 | 75 |
| | Nelspruit Location | 144 | 88 |
| | White River | 391 | 50 |
| | Nelspruit Town | 1,016 | 30 |
| | Karino | 460 | 49 |
| | Bosbokrand | 426 | 53 |

TABLE F: URINE EXAMINATIONS (NATIVE SCHOOL-GOING CHILDREN)

| Zone | Magisterial District | Number Examined | Percentage Positive | Zone | Magisterial District | Number Examined | Percentage Positive |
|----------|------------------------|-----------------|---------------------|---------|---------------------------|-----------------|---------------------|
| Zone I | Letaba | | | Zone II | Pietersburg | | |
| | Klein Letaba 6 schools | 321 | 35 | | Olifants 2 schools | 306 | 25 |
| | Moketsi 2 schools | 192 | 24 | | Cheunies 2 schools | 230 | 6 |
| | 1 school | 100 | 23 | | Marabastad 2 schools | 193 | 4 |
| | Letaba 10 schools | 876 | 50 | | Pietersburg 5 schools | 520 | 4 |
| | Olifants 4 schools | 273 | 53 | | North West 12 schools | 922 | 7 |
| Zone III | Waterberg | | | | Potgietersrust | | |
| | Sanddrift | 47 | 51 | | 7 schools | 813 | 13 |
| | Rooipoort | 101 | 27 | | Lydenburg | | |
| | Middelfontein | 101 | 50 | | (Highveld south of Steel- | | |
| | Rustenburg | | | | poort River) 3 schools | 218 | 19 |
| | Marikana, Tweedepoort | | | | Lowveld, north of Steel- | | |
| | and other schools | 1,178 | 33 | | poort River | 777 | 13 |
| | Groblersdal | | | Zone IV | Eastern Transvaal | | |
| | Dennilton | 283 | 32 | | Lowveld 11 schools | 700 | 56 |
| | Ottensville | 109 | 30 | | | | |

TABLE G: URINE EXAMINATIONS: EUROPEAN SCHOOL-CHILDREN

| Zone | Magisterial District | Number Examined | % Positive | Zone | Magisterial District | Number Examined | % Positive |
|------|--|-----------------|------------|------|----------------------|-----------------|------------|
| I | Letaba | | | II | Lydenburg | | |
| | Merensky School Farm | 295 | 10 | | Origstad | 130 | 17.4 |
| III | Groblersdal and surrounding areas | 668 | 11.2 | | Rustplaat | 32 | 3.1 |
| | Dennilton, Marble Hall, Groblersdal, Waterberg, Sine schools | 638 | 5 | | Three other schools | 183 | 5.2 |
| | Rustenburg (Schools 4) | | | | De Groot Boom | | |
| | Paul Kruger, Rustenburg, Kafferskraal, Elandsdraal | 1,563 | 4.4 | | Krugerpoort | | |
| | | | | | Burgersfort | | |
| IV | Eastern Transvaal | | | | Two schools (Komati- | | |
| | poort, Elandschoek) | 200 | 5.2 | | Barberton | 558 | 5.7 |
| | | | | | Nelspruit | 703 | 7.1 |
| | | | | | | | |

It is of marked practical importance to note that the distribution of bilharziasis is considerably more widespread than the distribution of the Bilharzia snail which is, after all, limited to the water sources or water holdings (dams, etc.) which are often a long distance apart. This means that in some areas, especially in the drier country to the west of the Drakensberge, the clearance of a relatively small number of streams and dams will prevent the transmission of bilharziasis over a large block of country.

DISCUSSION

Numerous problems and obstacles have been encountered. Most of these seem surmountable. Others will still have to be dealt with. It would be appropriate to cite just

a few of these obstacles and it will readily be appreciated that Bilharzia control is by no means as straightforward undertaking as was malaria control.

1. The success of any scheme for Bilharzia control must largely depend upon the willing co-operation of the farming community. In the Northern and North-Eastern Transvaal it has taken us many years to convince these farmers—individually, and through their various associations—that our mosquito larvicides are not harmful to their stock and their crops on which most of them are solely dependant for their existence. Sulphation offers the same problem and we expect the same or more difficulty in persuading the farmer that copper sulphate will have no deleterious effects. Fortunately we can give this assurance with some confidence as our biological and chemical tests

have shown (see back in this paper) how rapidly copper sulphate is rendered innocuous when added to natural collections of water.

2. During the rainy months (November to March) large numbers of snails are found in the grass pans which are such a distinctive feature of the Northern Transvaal. These snails are mostly *Bulinus tropicus*, but from 10% to 20% (see Table D) are the proven Bilharzia vector, *Physopsis africana*. Most of these pans dry up a few weeks to a few months after the last summer rains and may remain dry for many months before they are refilled with water. As soon as this happens the snail population re-appears. Where do these snails come from? We have found that these snails, with their openings sealed with epiphagms made largely of mud, aestivate in their thousands hidden in the depths of the dry tufts of grass and in between the grass roots. When placed in water the artificial epiphagm dissolves and normal life is resumed. We have raised new colonies of snails (*Bulinus tropicus*) from specimens taken from these pans as long as 18 months after the last water has dried up. Shousha similarly records that snails in Egypt and the Sudan will survive drying for 12 months, though apparently the mechanism is somewhat different. We have not yet found *Planorbis* to aestivate in this way, though it is possible that they do.

The resistance to drying naturally makes control more difficult as the aestivating snail is highly inaccessible to us and our molluscicides. It suggests that the best time for molluscicidal work in this type of water collection is when the water is at its highest as living snails do not appear to bury themselves when water is available.

3. Of at least as great significance as the incidence of Bilharzia amongst the human and cattle population is the incidence of bilharziasis amongst the natural snail population. This we have been unable to establish, as we cannot under field conditions differentiate between the various cercariae found, except to say that they are bifurcus or afurcus. De Meillon⁷ has previously commented on the difficulties found in identifying cercariae found under natural conditions on morphological grounds. It is considered though that for practical purposes, the recovery of typical forked-tail cercariae (either by placing the snail in fresh water—the ideal temperature is some 80° F though we have found shedding at as low as 50° F or by dissecting the snail) from known Bilharzia vectors (*Physopsis africana* and *Planorbis pfeifferi*) in an area where human bilharziasis occurs, is sufficient presumptive evidence to justify our considering these snails as dangerous and worth control. Any field work in this direction will, however, be of limited value until such time as all cercariae can be accurately identified.

It was a matter of regret to us that DDT—which we used extensively as a mosquito larvicide—is not sufficiently cercaricidal to be of value in the control of bilharziasis—as shown by Kuntz and Stirwalt.¹⁸ Incidentally, Jones and Brady¹⁹ showed that copper sulphate was only an effective cercaricide in concentrations above 50 p.p.m. and McMullin and Ingals²⁰ failed to kill the cercariae of *B. japonica* with concentration of copper arsenate, copper cyanamide or Paris green of up to 100 p.p.m. Gonzalez, Biaggi and Leon²¹ have, however, shown that a chlorine residual of 0.5 p.p.m. for 20 minutes will kill *B. mansoni* cercariae and this is of practical importance.

As yet the Department of Health has undertaken no systematic campaign to control bilharziasis and the work reported in this paper was conducted concurrently with the multitudinous other duties, especially malaria control, which keep the Malaria Unit of the Department busy in this part of the Union. It is hoped, nonetheless, that it will soon be possible to undertake a real control programme and in the meanwhile our preliminary work is continuing along the lines which appear to have the most immediate practical significance and in close liaison with biologists and all those interested in the preservation of our fauna.

It must be emphasized that the work was largely voluntary in nature and it was only the willing and enthusiastic team-work of all members of our unit—field staff, technical and administrative—which made it possible. To them we extend our sincere appreciation.

SUMMARY

1. Bilharziasis is known to occur in varying degrees of severity in the Northern and Eastern Transvaal—an area of over 60,000 square miles. The reasons are discussed why molluscicidal work is the only possible practical method by which the disease can be controlled in this area.

2. The distribution of the more important fresh-water snails is given for different parts of the Transvaal for the four quarters of 1950, and the distribution in various habitats is given for the known vectors *Physopsis africana* and *Planorbis pfeifferi* as they occurred during these four quarters. The close relationship between the occurrence of *Physopsis africana* and the incidence of *B. haematobium* is demonstrated and discussed.

3. Various molluscicides are discussed, and the opinion is given that copper sulphate is at present the molluscicide of choice (within economic limits). The instability of copper sulphate is demonstrated, and it is suggested how this chemical may best be used under our conditions.

4. Urine examinations done during the years 1937-1938 and 1948-1950 give percentage incidences of bilharziasis among Native school-children which vary from 88% to 4%, with the highest incidence in the low-lying country to the east of the Drakensberg mountains. The corresponding figures for European school-children vary from 17% to 3%. The patchy distribution is emphasized.

5. Problems created by the farming community's fear that copper sulphate may injure their stock or their crops is discussed.

6. The aestivation habits of certain snails are described, and a mention is made of the difficulty of identifying cercariae obtained from snails collected in their natural habitat.

7. It is indicated that this work has been of a preliminary and largely voluntary nature.

REFERENCES

1. Jopling, J. W. (1949): J. Trop. Med. Hyg., **52**, 121.
2. Mozley, A. (1944): *The Control of Bilharzia in Southern Rhodesia*. Rhodesian Printing and Publishing Co. Ltd.
3. Blair, D. M. (1948): S. Afr. Med. J., **22**, 462.
4. Shousha, A. T. (1949): Bull. World Health Org., **2**, 19.
5. Schwetz, J. (1949): Ann. Soc. Belge de Med. Trop., **29**, 37.
6. Porter, A. (1938): Publ. S. Afr. Inst. Med. Res., **8**, 1.
7. de Meillon, B. (1948): S. Afr. Med. J., **22**, 253.

8. Barlow, C. H. and Abdel Azim, M. (1948): 5th An. Rep. Bil. Snail Destruction Sec., Min. of Health, Gov. Press, Cairo.
9. Gillet, J. (1949): Ann. Soc. Belge de Med. Trop., **29**, 478.
10. Becker, J. G. (1916): S. Afr. Med. J., **11**, 156.
11. Khalil, M. and Halawani, A. (1947): J. Roy. Egypt Med. Assoc., **30**, 454.
12. da Mota, L. A. C. R. C. (1948): Ann. Inst. Med. Trop. Lisbon, **5**, 289.
13. McMullin, D. B. and Graham, O. H. (1947): Amer. J. Hyg., **45**, 274.
14. Nolan, M. O. and Berry, E. G. (1949): Pub. Health Rep., **64**, 942.
15. Nolan, M. O., Berry, E. G. and Gonzalez, J. O. (1950): Public Health Rep., **65**, 939.
16. Luthermoser, G. W. (1941): Rev. San. y Assist. Social., **6**, 874.
17. Bennie, I. (1949): S. Afr. Med. J., **23**, 97.
18. Kuntz, R. E. and Sturwalt, M. A. (1946): J. Parasitol., **32**, 529.
19. Jones, M. F. and Brady, F. J. (1947): Nat. Inst. Health Bull., **189**, 109.
20. McMullen, D. B. and Ingals, J. W. (1947): Amer. J. Hyg., **45**, 294.
21. Gonzalez, O. J., Biaggi, N. and Leon, R. J. (1945): Puerto Rico J. of Pub. Health and Trop. Med., **20**, 357.

ASPECTS OF CRANIO-CEREBRAL TRAUMA

I. THE SURGICAL ANATOMY AND PHYSIOLOGY OF CRANIO-CEREBRAL TRAUMA *

DAVID A. MUSKAT, CH.M.

General Hospital, Johannesburg

SURGICAL ANATOMY

A proper conception of the mechanics and pathological sequences of crano-cerebral injuries requires a knowledge of certain intracranial features. Trauma, especially through its widespread effects, can disorganize any and every part of the cranial contents, so that the anatomical configuration not only has an important bearing on the pathology of head injuries but may, and often does have a direct bearing on the death of the patient. Certain aspects of cranial anatomy therefore merit discussion.

The Skull. Except for certain thinned-out areas at the base, the inner and outer surfaces are condensed to form compact expansions of bone (the inner and outer tables). The outer table is thicker and roughened to allow for muscular and fascial attachments; the inner table is harder, approaching ivoryiness in parts, yet, for this very fact, possessed of a certain brittleness in structure. The diploic veins, in the soft, gritty and porous diploe between the tables, are wide, thin-walled and easily torn. Owing to adhesion to the surrounding diploe, they are unable to retract and so bleed profusely when torn. They lack valves and at intervals communicate with surface veins via the emissary veins, and with the underlying meningeal veins.

The bones of the vault are developed in membrane and retain a certain amount of elasticity; but the bones of the base, developed in cartilage, lack this feature, being more brittle. At places the bones of the base are paper-thin and may even be absent and replaced by a membranous layer of fibrous tissue (e.g. the cribriform plate).

The diploe has a certain amount of protective function, acting as a medium for the dissipation of forces applied to the exterior (*cf.* shatterproof glass—Dandy, 1947). However, more important protection is rendered by longitudinal and transverse thickenings of bone, giving it a buttress-like support. These occur at the glabella, the external angular process, the mastoid bones, the external occipital protuberance, the supraorbital ridges, the temporal crests, the occipital line, the longitudinal arch,

the thick posterior rim of the foramen magnum, the petrosal ridges and, more anteriorly, the sphenoid bones. Although these buttresses are thickened parts of the bony structure, they exhibit a tendency to brittleness, accounting for the frequency with which fracture-lines cross them. It is more usual, however, for fissures to be deflected and deviated to points of weakness such as foramina, the cribriform plates, the pituitary fossa, the lateral parts of the petrous bone, the sphenoidal fissure, the roofs of the orbits and the tegmen tympani (Rowbotham, 1949).

In the infant the vault of the skull is large compared to the base and the membranous layer between the sutures is of considerable thickness. The mobility of the bony plates and the fibrous attachments of the vault tend to confine the effect of violence to the bones primarily struck, with but little transmission of the force to other parts of the cranium (Vance, 1927). This very fact, however, renders the underlying soft brain more vulnerable to injury. With increasing age sutural lines are lost, the skull bones become more brittle and thicker, mainly due to deposition of bone on the inner table to replace the shrinkage of the brain.

It is generally taught that the Bantu skull demonstrates differences in texture and thickness from that of the average European. Certain craniological differences undoubtedly occur. However, a small sphere exhibits greater solidarity than a large one, and some of the alleged increased resistance of the Bantu skull to force, may be due to the fact that the average weight of the European brain is 1,475 gm. compared to that of 1,331 gm. in the Negro (Weaver, 1930). Personal experience in operating on the Bantu skull, observation of skulls at autopsy, perusal of autopsy records and the observation of experienced pathologists, have failed to justify the conviction that the skull of the average Bantu is thicker or tougher than that of the average European.

The Dura Mater. This thick, resistant, inelastic membrane forms a lining for the bony cavity and a mechanical support for the brain and the large venous sinuses. It has the character and functions of a fibroblastic membrane (Leary and Edwards, 1933). The outer layer is firmly adherent to the skull, particularly in the very young (*cf.*

* The References will be published at the end of the concluding article in this series.



Easing the way ...



Although the adage 'life begins at forty' may be true in theory, it is in practice that we realise that it is not long before the difficult milestone of the menopause is reached. The years of stress may be eased by the timely administration of a preparation designed to counteract the depression, nervous phenomena, and vasomotor disturbances so troublesome to women patients.

Euvalerol M contains an odourless preparation of valerian with $\frac{1}{4}$ grain (16 mg.) phenobarbitone and 0.1 mg. stilbæstrol in each fluid drachm. Its use is followed by marked diminution of symptoms and rapid restoration of emotional balance.

EUVALEROL M

In bottles of 8 fluid ounces.



Literature on application.

ALLEN & HANBURY'S (AFRICA) LTD
(INCORPORATED IN ENGLAND)
409-11 - SMITH STREET - DURBAN

(EMF, 2)

Make these seven important urine tests in one minute with DROP TEST in your consulting room

| | | | | | | |
|----|---------|-------|---------|---------|------|--------------|
| pH | Albumin | Sugar | Acetone | Indican | Bile | Occult Blood |
|----|---------|-------|---------|---------|------|--------------|

DROP TEST is the *only* kit available that permits you to make a *complete* and clinically accurate on-the-spot chemical examination of urine for pH, albumin, sugar, acetone, indican, bile and occult blood. The entire urinalysis is completed in one minute—a drop of urine and a drop of reagent. Results are easy to read and as precise as conventional methods.

DROP TEST reagents cost much less than standard reagents, but DROP TEST's greatest economy is in the 20 to 1 saving of technicians' time. The kit is unconditionally guaranteed for one full year of stability and accuracy. DROP TEST carries the seal of acceptance of the American Medical Association. Detailed literature is available, without cost or obligation.



Manufactured in U.S.A. by Biochemical Methods, Inc. Sole distributors for South Africa and the Rhodesias:

PETERSEN, LIMITED

P.O. BOX 38, CAPE TOWN

P.O. BOX 5992, JOHANNESBURG



Oxyenaire
South Africa (Pty.) Ltd.

THE "OXYAIR" FACE MASK

THE "OXYAIR" FACE MASK, WHICH HAS BEEN INTRODUCED INTO THE FIELD OF OXYGEN THERAPY BY OXYENAIRE, HAS A THIN MOULDED FACEPIECE OF TRANSLUCENT PLASTIC. IT INCORPORATES A GAUZE VENTILATION HOLE, RESERVOIR BAG, A SMALL BORE PLASTIC TUBE FOR OXYGEN SUPPLY, AND THE WHOLE HAS ELASTIC SUSPENSION.

THE GAUZE-COVERED VENTILATION HOLE PROVIDES A MINIMUM RESISTANCE TO BREATHING AND ALSO CREATES A TURBULENCE WHICH IS AN ADVANTAGE AT HIGH INSPIRATORY AND EXPIRATORY FLOW. THE CONNECTION OF THE RESERVOIR BAG ON TO THE MASK IS SO DESIGNED AS TO AVOID ANY POSSIBILITY OF CLOSURE OF THE NECK OF THE BAG WHEN THE PATIENT IS IN A RECLINING POSITION. SHOULD THE OXYGEN SUPPLY FAIL, THE LOW RESISTANCE OF THE VENTILATION HOLE ENSURES ADEQUATE SUPPLY OF AIR. THE OVERALL WEIGHT OF THE MASK IS 2 OZ.

FURTHER PARTICULARS MAY BE OBTAINED FROM:

Reg. Off. : 41 Main House, Main Street, JOHANNESBURG
P.O. Box 770 Telephone 33-1137

MIXED DIET MENUS AND RECIPES

NUTRINE FEEDING

It should be realised that no hard and fast rule can be laid down as to the exact age when the Mixed Diet should be introduced.

The following menus are for babies of average age and weight, and may require considerable adjustment for backward or particularly advanced babies. Here you are asked to make use of the services of our Free Advice Bureau if advice is not available.

Bear in mind that before new foods are introduced the artificial should be well established on the particular food mixture which is being given. The three-day plan is an excellent one when making changes in food. This gives the mother time to judge whether any particular item is accepted.

If baby cannot tolerate pure COD LIVER OIL, suitable substitutes are recommended in our Mothers Nutrine Guide.

MENU FOR BABIES 5 MONTHS OLD.

| 6 a.m. | 10 a.m. | 2 p.m. | 6 p.m. |
|---|---|--|--|
| Full Nutrine Feed, 8 ozs (see end of menu). | Cod Liver Oil (15 drops). A taste of coddled egg yolk on a finger of white, baked crust of rusk, twice during the first week, three times the second week, and thereafter daily. When eggs are very expensive the finger of rusk can be given or crust can be given 10 minutes before this feed every day, and ½ teaspoon of raw egg yolk added to the bottle feed, as frequently as recommended for the coddled egg. | Cod Liver Oil (15 drops). After Baby has had the egg yolk at 10 a.m. for 1 week, strained vegetables can be given about 15 minutes before the 2 p.m. bottle feed, on the days when egg yolk is not given. (Follow recipe for Vegetable Broth on page 6.) Give Baby 3 or 4 teaspoons of the broth 15 minutes before the 2 p.m. Nutrine Feed. Much may be wasted at first, therefore the full bottle feed may be given. After the first week, the vegetables through the sieve and increase gradually to 2 ozs. The broth can then be given daily, or as a change, and to save labour, give a little Marmite in the proportion of ½ teaspoon to 2 ozs. of warm boiled water. | Cod Liver Oil (15 drops). After Baby has had the egg yolk at 10 a.m. for 1 week, strained vegetables can be given about 15 minutes before the 2 p.m. bottle feed, on the days when egg yolk is not given. (Follow recipe for Vegetable Broth on page 6.) Give Baby 3 or 4 teaspoons of the broth 15 minutes before the 2 p.m. Nutrine Feed. Much may be wasted at first, therefore the full bottle feed may be given. After the first week, the vegetables through the sieve and increase gradually to 2 ozs. The broth can then be given daily, or as a change, and to save labour, give a little Marmite in the proportion of ½ teaspoon to 2 ozs. of warm boiled water. |

Nutrine Mixture for the Day.—Take the quantities for 5 months on the blue label or red label tins.

offered to doctors . . .

Nutrine is prescribed in almost every instance as a suitable food to be included in the diet of babies from 4 to 5 months old. For this reason the manufacturers, Hind Bros., have prepared a pamphlet on Mixed Diet for use in conjunction with Nutrine feeding and now offer it to Doctors in the hope that it will prove helpful to them and to their patients.

MIXED DIET PAMPHLET

To "Mixed Feeding," Dept. 85, Hind Bros. & Co. Ltd., Umbilo, Natal. Please send me a supply of Mixed Diet Menus and Recipes in English/Afrikaans.
(Strike out language not applicable.)

NAME

ADDRESS


NUTRINE

is Nearest to Nature

In the treatment of—
**ANGINA PECTORIS, BRONCHIAL ASTHMA,
CORONARY THROMBOSIS, CHRONIC COR PULMONALE**

A highly purified form of **SHELLIN**, the active principle obtained from the seeds of the Eastern Mediterranean plant *Amor-vivaga* (Liliaceae).

Benecardin
REGD.



Produces a rapid yet prolonged action
without lowering the blood pressure.

Literature on request to:

BRITISH CHEMICALS & BIOLOGICALS (S.A.) (PTY.) LIMITED

P.O. BOX 5788 259 COMMISSIONER STREET, JOHANNESBURG

the rarity of extradural haematoma at this age); adhesion diminishes with age but in the very old it again tends to be more marked. Through the various adhesions pass minute meningeal arterial twigs and fibrous processes which tear in the process of separation of the dura from the inner table.

Of the various dural septa the *tentorium cerebelli* assumes important significance in crano-cerebral pathology. The concave free edge, anteriorly, forms the margin of a space—the incisura tentorii—in which lie the midbrain, the basilar vessels, the posterior cerebral vessels, the oculomotor nerves and the anterior part of the superior vermis of the cerebellum. The space of Bichat, between the mid-brain and the sharp crescentic border, is minimal, allowing for the free flow of cerebrospinal fluid between the infratentorial and the supratentorial cisterns. A firm, resistant structure thus effectively divides two important regions of the brain from each other, viz. the infratentorial compartment containing the 'vital structures' (medulla, pons, cerebellum) from the supratentorial compartment with its contained cerebral hemispheres, a relatively narrow, soft core of brain tissue intervening between the two.

The dura mater is richly supplied with arteries and veins, particularly in the fossae. Anteriorly it is supplied by meningeal branches of the ethmoidal and internal carotid arteries. In the middle cranial fossa are the middle and accessory meningeal branches of the internal maxillary artery, and branches of the ascending pharyngeal and internal carotid arteries. More posteriorly we find branches of the occipital, and anterior and posterior meningeal branches of the vertebral arteries. The great vascularity of this dense, adherent membrane is one of the major contributing reasons for the gradually or rapidly increasing space-occupying lesions created by haemorrhage, the ever-increasing haematoma breaking down more and more communicating twigs, giving rise to what one might appropriately call a vicious expansion. The dura is particularly adherent in the region of the cribriform plate, where it is also thinnest. It is, moreover, very firmly anchored by its extensions through the bone along the olfactory nerves (Cairns, 1937).

The *subdural space* can be likened to the peritoneal and pleural spaces, where the smooth surfaces are separated by a thin film of moisture to allow for movements of organs; it must be remembered that the brain is a constantly moving (i.e. pulsatile) organ. The space is potential, being in life merely a cleft. Embryologically the formation of the space probably derives from the separation of the ectodermal pia-arachnoid from the mesenchymal inner layer of the dura mater: hence the difference in properties of these membranes. The pia-arachnoid is poorly vascularized and has weak fibroblastic properties, whereas the dura reacts strongly to infection and trauma and can organize and remove blood from the subdural space (Leary and Edwards, 1933).

Certain Important Vessels. The anterior branch of the middle meningeal artery crosses the great wing of the sphenoid and in its anterior inferior angle, with its *venae comites*, grooves the bone deeply or may even be completely enclosed in a bony tunnel for a part of its course. It and its main branch follow roughly the line of the precentral sulcus. The artery is closely adherent to the

dura, particularly where it is semi- or completely encased by bone (in 60% of cases—Gurdjian and Webster, 1942) so that it is almost impossible to separate the dura from the skull without causing rupture of the artery or its companionate veins, the thin-walled nature of the latter facilitating injury, with resultant extradural or even subdural haemorrhage.

The *posterior cerebral arteries* are the terminal branches of the basilar artery. Each artery passes lateral to and immediately above the oculomotor nerve, where it lies in very close relation to the free edge of the tentorium cerebelli. Winding round the cerebral peduncle it reaches the under surface of the cerebrum to supply the temporal and occipital lobes. From its position between the mid-brain and the third nerve medially and the free edge of the tentorium laterally, it can readily be understood how a supratentorial herniation of the uncus can squeeze and even occlude the artery with resultant ischaemia or infarction of the area of brain supplied by it (including the cortical visual areas).

The Cortical Veins. Of surgical importance are the superior cerebral veins draining the supero-lateral and medial surfaces of each hemisphere. The anterior veins drain into the longitudinal sinus almost at right angles, whilst the larger posterior ones run obliquely forwards, upwards and medially, thus entering the sinus in a direction opposite to the flow of blood inside it. The relatively unsupported veins are thin walled and elastic, permitting participation in the normal variations of intracranial pressure. They leave their partially protected beds in the sulci to enter the sinuses by piercing the arachnoid mater and traversing the subdural space for a short but variable distance, finally to enter the longitudinal sinus, a relatively fixed point. The transition between a potentially moving cortical surface to a fixed dural point is an important aetiological factor in the causation of subdural haematomata.

The *uncus* is an important portion of brain tissue, at times playing a major role in abnormal intracranial hydrodynamics. It is the anterior recurved end of the hippocampal gyrus, lying in juxtaposition to the third nerve and in close relation to the anterior end of the tentorium cerebelli, and abutting on the side of the mid-brain. In conditions of increased intracranial pressure above the tentorium, the uncus, from its position, is forced downwards into the space of Bichat, nipping the third nerve between the midbrain and tentorium, and is responsible for certain important clinico-pathological consequences.

SURGICAL PHYSIOLOGY

1. The Cerebrospinal Fluid. The inferior surfaces of the frontal and temporal lobes (and the tips of their poles) are separated from the floor of the skull by only a thin layer of intervening fluid. Absence of cerebrospinal fluid cushioning is one of the factors predisposing these regions to the direct and indirect effects of trauma. Although it is generally accepted that absorption of fluid is effected by the arachnoidal villi and granulations, there are some who believe that there is a direct passage of cerebrospinal fluid into the blood stream (Howe, 1928; Dandy, 1947), probably by osmosis. A major function of the cerebrospinal fluid is its cushioning or dampening effect in

protecting the brain from external jars and forces. In order to maintain this function a certain amount of pressure is required.

2. *The Intracranial Pressure.* It has long been taught (Monroe-Kellie doctrine) that the total volume of the cranial contents is fixed by the rigid confines of the bony walls. In order to maintain pressure equilibrium, a change in any one component should be associated with a reciprocal change in some other, i.e. brain, cerebrospinal fluid or blood. In contrast to this older conception of the closed box, the cranio-vertebral cavity is now recognized to have a certain amount of elasticity, due particularly to the remarkable resiliency of the cerebral circulation (Loman, 1938). One can regard the cranio-vertebral contents as comprising two types of elements.

1. Static or inelastic;

2. Dynamic or elastic (mainly the vessels, particularly the thin-walled veins).

The normal limits of the cerebrospinal fluid pressure range from 50-200 mm. water, i.e. with the patient in the lateral horizontal position. In the erect or non-horizontal position the pressure is not so much related to the column of cerebrospinal fluid above the needle as to the hydrostatic and elastic changes occurring in the venous wall surrounding the fluid, e.g. upward tilting causes a lowering of the venous and hence of the cerebrospinal fluid pressure (Loman). In the erect position the normal adult intracranial pressure varies from 50-150 mm. water below atmospheric pressure. The zero point is about the level of the lower cervical vertebrae (Munro, 1938).

3. *The Role of Vascular Factors.* Opinion favours a sympathetic nervous control of the cerebral blood vessels. In addition, contraction of the smaller vessels can occur independently of the nervous stimulation, as, e.g. proved by Echlin (1942) who showed that pial vessels can contract vigorously locally when stimulated under experimental conditions. The cerebrospinal fluid pressure is closely related to the venous pressure and the cerebral blood flow to the arterial pressure, the combination of these two factors tending to keep oxygenation of the brain at a uniform level. Paradoxically, when the systemic blood pressure falls there occurs a marked vasodilatation, the critical level at which this occurs being about 70-80 mm. Hg (Howe, 1928). The vasodilatation in the face of a falling blood pressure can actually provide a greater blood flow, provided the fall is not too great, e.g. by histamine or amyl-nitrate (Loman). CO₂ is a potent vasodilator, and alcohol to a lesser extent, but as a result of the concomitant general vasodilatation, the blood pressure may be sufficiently lowered to cause a decrease in the cerebral blood flow. Vasodilatation can also occur reflexly in concussion by reflex stimulation of the vagus nerves (Denny, Brown and Russel, 1941), probably via the parasympathetic fibres of the seventh cranial nerves (Forbes, 1940). Scheinker (1944) considers vasoparalysis a frequent reaction of the blood vessels of the nervous system, caused by acute trauma or intoxication. This circulatory stasis or vascular damage, once established, would result in progressive cerebral ischaemia, anoxia and oedema (Masserman, 1934).

Oedema tends to reduce the cerebral blood flow by compression-narrowing of the capillaries and venules. Loss of fluid from the perivascular spaces, on the other

hand, tends to dilate the capillaries and veins but, though increasing the amount of blood in the vessels, will not necessarily increase the flow, particularly when, as in these cases there is often a generalized loss of fluid, causing not only a lowering of systemic blood pressure but also an increased viscosity and resultant peripheral resistance (Forbes, 1940). Cerebral blood flow is also reduced by a rise in the general intracranial pressure above 300 mm. water (Wright, 1945). Adequate blood flow is important as the brain has a high level of metabolism, causing venous blood leaving the brain to be extensively reduced. Energy utilization in the brain is entirely provided by carbohydrates and it is here that the role of vitamins, particularly the B group, is so important. These factors influence O₂ consumption greatly and therefore are intimately related to the blood flow.

Any condition causing compression of the cerebral veins or impeding the venous outflow from the brain will cause a rise in intracranial pressure. This occurs, e.g. in the well-known Queckenstedt test, also in the head-down position and conditions increasing the intrathoracic pressure, e.g. coughing, straining at stool, laughing, etc. A generalized rise in intracranial pressure results in the following sequence of events (Forbes, 1940): The compression of the veins results in a slowing of the rate of blood flow and a decrease in the volume of output. Venous stasis follows and capillary pressure rises sufficiently to overcome the obstruction and the circulation is resumed. The venous and capillary pressures can rise greatly without approaching that of the larger arteries. However, if this compensatory force rises high enough to approach the level of the arterial pressure, a rise in the general arterial pressure will occur reflexly. Gradual or moderate increases in intracranial pressure may not interfere with the cerebral flow so long as the arterial pressure remains sufficiently high. To quote Loman: 'It is only when the brain is suddenly overwhelmed by great increases in intracranial pressure that the systemic arterial pressure responds to aid the flow of cerebral blood.'

4. *Intracranial Tension and the Classical Syndrome of Cerebral Compression.* The names of Kocher and Cushing are closely associated with the compression syndrome. The former, in his postulates, has set forth the sequence of events occurring as the result of increasing intracranial tension. He divided the course of compression into a series of stages:

Stage 1. A mild compression encroaches mainly on the venous bed, causing slight congestion and increase in capillary pressure.

Stage 2. Venous and capillary stasis are further increased, there is a relative anoxia of the vital bulbar centres, reflexly eliciting a slight rise in the systemic blood pressure, and a slight slowing of the pulse rate. Early symptoms of compression become manifest: headache, excitement, stupor, possibly focal signs.

Stage 3. Advancing compression results in capillary anaemia. The medullary centres attempt to react by a marked increase in blood pressure and slowing of the pulse. Advanced symptoms now manifest themselves: respiratory changes, e.g. Cheyne-Stokes breathing, coma, pupillary and optic disc changes. The prognosis, previously good, now becomes grave.

Stage 4. Increasing pressure fails to elicit medullary reflex responses, i.e. the medullary centres become exhausted. The blood pressure falls, there is marked increase in the heart rate, the pupils become dilated and fixed, the respiration slows down

and finally ceases. This is the stage of decompensation and the prognosis is hopeless.

The above sequence of events has been accepted universally, though the fact that large numbers of cases of cerebral compression do not exhibit the classical syndrome demonstrates that it is unwise, even dangerous, to make the above teaching a dogma. When present, the signs are certainly important guides to the clinical assessment of the case, but their absence does not rule out a vicious expanding lesion. Experiments carried out by the writer have failed to confirm the classical march of events in compression.

Intracranial pressure can be increased either by subjecting the normal quantity of cerebrospinal fluid to increased tension (as by a space-occupying lesion); or by increasing the amount of fluid. The latter can be effected either by diminishing the rate of absorption, increasing the rate of secretion or by bleeding into the subarachnoid space. In addition, haemorrhage can interfere with the absorption of cerebrospinal fluid in several ways:

1. By mechanically blocking the arachnoid villi.
2. As a result of contusion and laceration of the brain there is a loss of absorption area for the pachionian corpuscles (Shatara, 1936).
3. The communications between the intracerebral and extracerebral cerebrospinal fluid spaces are small and easily blocked and it is reasonable to assume, especially if coagulation occurs, that this may result in cutting off cerebrospinal fluid from the absorptive area.

Venous engorgement, a frequent accompaniment of brain swelling, results in increased intravenous pressure and interferes with normal fluid absorption, thus causing increased cerebrospinal fluid pressure. It is thought by some (Scott, 1940; Rand and Courville, 1931) that trauma may increase secretion due to abnormal activity or irritation of the choroid plexuses and ependyma.

Cushing showed experimentally that when the intracranial tension reached that of the arteries, the vasomotor centre responded reflexly, eliciting a rise in blood pressure, often to twice its normal level. Eyster (1901) proved that intracranial pressure of less than the mean blood pressure (90-100 mm. Hg) had very little effect upon blood pressure, pulse or respiration. Pressures encountered in trauma are rarely, if ever, capable of setting in motion the train of physiological events leading to medullary paralysis. Browder and Meyers (1936, 1938) find that a rapidly increasing intracranial pressure with its 'compensatory' syndrome has no application in clinical cases; and the reverse, viz. a compensatory syndrome, does not indicate increased cerebrospinal fluid pressure. In their experience repeated observations on the pressure cannot by themselves be employed as an index of the clinical condition of the patient, of his likely course, or of the necessity of surgical intervention. Such also must be the experience of most who are closely associated with this form of trauma. The conception of the classical compression syndrome so long taught can no longer be accepted in its entirety. Investigation of the course of severe and fatal head injuries have not upheld its tenets. There are indications that it is highly probable that many patients are losing their lives because their case sheets do not fall in line with the classical picture.

Zierold (1935) observed that when an abnormal intra-

cranial pressure did occur, it was the result of haemorrhage, either extradural, subdural, subarachnoid or intracerebral. Crawford goes so far as to say that intracranial tension is not the basic problem and Russell (1932) found no evidence that increased pressure contributed to a fatal issue; low pressures are often found in the terminal stages of head injuries. Frequently successive daily punctures reveal pressures tending to normal when actually the patient was becoming comatose and operation demonstrated haematomata under quite considerable pressure (Brody, 1940). Incisural or foraminal blocks probably play important roles in these cases. Furthermore, a normal reading in a dehydrated patient (as these patients so often are) may be misleading, as the actual pressure is probably very much higher (Munro, 1938).

5. *The State of Consciousness and Intracranial Pressure.* One school of clinical observers thinks that unconsciousness *per se* is a most important gauge of the degree of intracranial pressure. Cannon, Scott, Gurdjian and Webster, to mention a few, experimentally or clinically proved high pressures to be related to unconsciousness. Denny-Brown and Russell (1941), however, found no evidence that increased pressure contributed to a fatal issue or that there was any relationship between the degree of unconsciousness and the spinal pressure. Zierold (1935) and Browder and Meyers (1938) likewise found no relationship between unconsciousness and pressure. Rowbotham (1949) found a moderately raised pressure in the majority of his unconscious patients. It is the writer's opinion, from an analysis of over 100 fatal cases, that the state of consciousness has no relationship to the degree of intracranial pressure as measured by lumbar manometry, but this does not mean that there is no raised intracranial pressure or that some remediable space-occupying lesion is not present—a fact that cannot be sufficiently emphasized.

Whether unconsciousness is due to disturbed thalamic function (Greenfield, 1938), frontal lobe function (Dandy), changes in the metabolism of neuronal cells (Walker, Kollross and Case), related to direct traumatic paralysis of the cortical neurones (Eden and Turner), anaemia of the brain (Trotter, Cushing, Scott), or injury to subcortical structures and the midbrain (Jefferson), there is no doubt of its profound bearing on the ultimate prognosis of the case. It is a most important indication when prolonged (12 hours or more), of the secondary effects of head trauma.

Loss of consciousness has always found its classical application in the development of the 'compression' syndrome. The lucid interval is the well-known feature of this development. The essential condition is the occurrence of a second phase of disturbance of consciousness, in which, owing to the gradual, or rapid formation of a space-occupying lesion, the patient lapses into the coma of compression, as distinguished from the unconsciousness of concussion. However, so frequently is the primary phase associated with severe disturbances of consciousness, that the secondary phase is a mere continuation, or superimposition, and no 'interval syndrome' is manifest. It is for this reason that any case showing prolonged unconsciousness must be regarded as due to a space-occupying lesion until proved otherwise by diagnostic craniotomy, and the diagnosis of the primary injury only verified by ruling out the secondary complication. The

second and third days constitute the critical period for unconscious patients (Hawkes, 1943). It is probably due to the fact that in this group the secondary manifestations are of rapid progress and contain the operable cases, in which failure to remove the cause of the compression, theoretically at any rate, inevitably leads to a fatal outcome. Munro goes so far as to state that no head injury should die after the third day!

6. *The Blood Pressure.* Browder and Meyers (1938) dispute the accepted dictum that increasing intracranial pressure produces its vital effects through stimulation of the medullary centres. Although experimentally the cerebrospinal fluid pressure must equal the systolic blood pressure before a rise in systemic blood pressure results, the clinical value of a change in blood pressure, according to these authors, was insignificant and possibly misleading. Dandy, Zierold, Eyster and Fremont-Smith and Merritt likewise found the blood pressure of little value in head injuries. Others, however (Denny-Brown and Russell, Walker *et al.*, Gurdjian and Webster, Pearce Baily) attach considerable importance to a rise in blood pressure, though Gurdjian (1933) himself found the pressure determination to be valueless as an index of increased intracranial pressure. It is probable, as stressed by Rowbotham (1949), that a rise in systolic pressure not infrequently occurs, often associated with a full, bounding pulse (due to the increase in pulse pressure), giving it the so-called 'vagal quality'. However, this rise is not always associated with a 'demonstrable rise in intracranial pressure and some other mechanism, in the nature of a vago-glossopharyngeal reflex, may play a part, possibly from the concussional injury.

7. *The Pulse.* The importance of the pulse rate has long been impressed on medical students and the so-called characteristic slowing has almost been accepted as a medical axiom—it is regarded as almost a travesty of medical supervision to omit the 'hourly pulse rate' chart. That occasionally a slowing of the pulse rate indicates exactly what it is meant to convey is not denied, yet not only can it be a false alarm, but the fact that this slowing has not occurred has been the reason for not suspecting increasing mischief of an operable nature and procrastination had inevitably led to verification on the post-mortem table or, too late, on the operating table. Of course, reliance on one sign alone in such a case is not good surgical practice, but it supersedes, in the mind of the practitioner, all the other signs of the 'compression syndrome', and it is rarely that one finds hourly recordings of blood pressure, temperature, respiration, states of consciousness, etc., all presumably showing characteristic variations as well. Kocher's postulates seems to have made of the pulse rate almost a mathematical formula, i.e. its variation inverse ratio to the intracranial tension. So many factors influence the circulatory system, however, that neither the pulse nor the blood pressure react to intracranial conditions with any degree of consistency (Vance, 1927). Marked bradycardia, apparently due to vagus excitation, can occur in simple concussion (Jefferson, 1938; Bagley, 1929). The occurrence, however, of a slow pulse in the stuporose or comatose patient is of important clinical significance; bradycardia, to be of any prognostic value, must always be considered in conjunction with the state of conscious-

ness. Analysis of hourly pulse charts in a large series of fatal cases reveals practically always inconsistencies, the record showing a rate e.g. of 72, an hour later 64, then 80, 88, dropping down to the 60's and rising and falling irregularly until the terminal 'racing pulse' supervenes. Except for a few isolated instances, the pulse rate cannot be regarded as a reliable therapeutic index of intracranial traumatic pathology.

8. *The Respiration.* Classically, as in the case of the vasomotor centre, the respiratory centre responds to an increase in intracranial pressure (to the level of that in the arteries), manifesting itself in the well-known Cheyne-Stokes phenomenon. The respiratory centre, however, is more vulnerable and death in cranio-cerebral trauma is almost invariably ushered in by respiratory failure. The stertorous breathing seen in the severe head injury may be due to vagal inhibition or merely the generalized loss of tone found in all deeply unconscious patients. Typical Cheyne-Stokes respiration is unusual, but as the brain stem is a very vulnerable part in severe injuries, as manifested by the frequent occurrence of haemorrhages, it is not inconceivable that the direct effect of trauma on the respiratory centre either macroscopic, microscopic or neuronal, may play an important part in causing any type of respiratory variation and leading to a fatal issue.

9. *The Temperature.* Mild and moderate injuries are not characterized by any special disturbance in the temperature, which may rise to 100°-101° F. A rise in temperature has been one of the accepted findings in increasing intracranial pressure, similar to the other reactions of the vital centres, so much so that Stevenson, e.g. advocated 15-minute temperature readings, anything above 103° F being regarded as evidence of increasing intracranial damage from compressional effects. Blood in the cerebrospinal fluid may cause the temperature to rise to 102° F or even 103° F due to a 'sterile meningitis' or absorption of broken-down blood proteins; but when there is an early rise to this level or when it exceeds 103° F, it is indicative of serious interference with the thermoregulatory centres.

Classically hyperthermia is attributed to haemorrhages in the pons (along with 'pin-point' pupils) and Courville (1945) considered focal haemorrhages, with or without oedema, in the midbrain or pons, to be frequent causative factors. By following up the clinical records of patients exhibiting these haemorrhages *post mortem*, the writer has not been able to verify these generally accepted dicta and there has been no correlation between pontine haemorrhages and the alleged hyperpyrexia.

Vomiting. Occurring early, this is usually due to recovery from the concussive phase, to the gastric irritation of alcohol or, in some cases, from ingested blood. In the later phase of cranio-cerebral trauma it may be significant, yet most authorities (e.g. Zierold; Browder and Meyers) disregard its importance as evidence of raised intracranial pressure. Vomiting in the deeply unconscious patient, however, is gravely ominous, often enough preceding death.

10. *The Reflexes.* Two series of reflex changes occur:

1. Those related to the actual concussive phase;
2. Those associated with the development of secondary complications.

In regard to the former, loss of the corneal reflex is

PACYL

mm Hg

220

210


200

190

The sovereign treatment of essential,
climacteric or arteriosclerotic

HYPERTENSION

and its
concomitant symptoms



Relief is also afforded to those patients in whom sclerosis is so far advanced as to make any actual reduction in blood pressure impossible, and to those who suffer from vasomotor disorders, which are not accompanied by raised blood pressure.

Supplied in bottles of 50 and 200 tablets.

BASE: A choline derivative, originated and developed in our laboratories.

Woman, age 50, with hypertension.
F. H. LEWY, Z. Klin. Med.
107/1-2.

60 clinical papers have confirmed the therapeutic value of Pacyl, which was awarded the Diploma and Gold Medal at the 7th International Congress of Hygiene, 1933.

VERITAS DRUG COMPANY LIMITED

LONDON AND SHREWSBURY - ENGLAND

For further information and samples apply to our Distributors in South Africa:

LENNON LIMITED - P.O. Box 8389 - JOHANNESBURG

APONDON

For **SAFE** weight reduction
by pharmacologically
DETOXIFIED Thyroid

Announcing

PENDEX

**a penicillin-vasoconstrictor combination
for intranasal use**

• 'Pendex' answers the demand for an aqueous penicillin-vasoconstrictor for intranasal use. It presents, in a stable aqueous solution, the potent antibacterial action of penicillin combined with the rapid and prolonged vasoconstriction of 'Paredrinex'. An important advantage of this new form of penicillin is its superior stability. The value and clinical applications of 'Pendex' will be immediately apparent. Detailed information about this striking advance in penicillin therapy will be sent on request.

Available on prescription only in 15 c.c. bottles with dropper.

When prepared as directed 'Pendex' will contain for one week at room temperature, not less than: Crystalline potassium penicillin G, 1,500 International Units per c.c.; 'Paredrinex', 1 per cent.; in a specially buffered aqueous solution.



The 'Pendex' package consists of penicillin in a dry state and a buffered aqueous solution of 'Paredrinex' — each in a separate container. The pharmacist has only to mix the two, and 'Pendex' is dispensed freshly prepared and with the penicillin at full therapeutic potency.

PHARMACAL PRODUCTS (PTY.), LTD., 2 DIESEL STREET, PORT ELIZABETH

for Smith Kline & French International Co., owner of the trade marks 'Pendex' and 'Paredrinex'
Distributors in Rhodesia: Geddes Ltd., P.O. Box 877, Bulawayo

significant of a sensori-motor disturbance of the medulla, occurring at the time of the acute traumatic disturbance of cerebral function. During the deep coma of primary injury the limbs are generally completely flaccid and all the superficial reflexes are usually abolished. A characteristic feature, however, is the variability of the reflexes, including the plantar reflex, so much so that they have been regarded as being of no significance (MacGregor, 1942).

When complications, particularly haemorrhage, become superimposed, clinically very valuable alterations may occur. The manifestation of unilateral hyper-reflexia is very often indicative of the development of local compression, especially when associated with significant pupillary changes. In this respect it is the unilateral and, in the late case, bilateral fixed dilatation of the pupils that is the most important change. The mechanism is probably as follows: As the uncus descends in a supratentorial herniation, it compresses the third nerve or stretches it (with the slight downward displacement of the midbrain). The effect on this nerve, containing the parasympathetic fibres of the sphincter mechanism of the pupil, is at first to stimulate it, causing a slight transient phase of pupillary constriction. Increasing pressure paralyses the sensitive parasympathetic fibres, leading to sympathetic overaction and a gradual dilatation of the pupil. During this phase there may be a sluggish reaction to light. Finally the pupil becomes fully dilated and fixed. By this time the brain stem is being compressed against the opposite edge of the incisura tentorii (or the opposite uncus commences to herniate), giving rise to a similar chain of effects until finally both pupils are widely dilated and completely immobile. Intrinsic primary or secondary changes in the brain stem may play a part in disturbing the oculomotor mechanism. In some late cases, complete third nerve paralysis may occur, resulting in partial ophthalmoplegia and ptosis. An expanding lesion in the sphenotemporal region is particularly dangerous in initiating uncal herniation, usually on the same side, and thus causing an ipsilateral fixed dilated pupil, indicating the site for initial exploration. It is not unusual to find spasticity and hyper-reflexia on the same side as the dilated pupil. In these circumstances it is postulated that the brain stem is compressed against the opposite sharp edge of the incisura, causing a lesion of the opposite pyramidal tracts and thus ipsilateral reflex and other changes; hence the advice to explore first on the side of the dilated pupil.

11. *Anoxia and its Effects on the Brain.* The brain is extremely avaricious for O_2 , particularly the grey matter,

which utilizes four times as much as the white (accounting for the greatly increased capillary bed in the former). It can only use up energy derived from carbohydrates, principally in the form of glucose, this substance and O_2 being taken up in approximately equal amounts. By phosphorylation glucose is transformed, in a series of steps, ultimately to pyruvic acid, and this again, in an adequate supply of O_2 , to CO_2 and water. If there is insufficient O_2 , the process is incomplete and pyruvic acid is converted to lactic acid. Cerebral anoxia has been shown to cause an increase in the cerebral lactic acid; likewise thiamine deficiency and, to a certain extent, most anaesthetics. There is now sufficient evidence to show that anoxia plays an integral part in the physio-pathology of cranio-cerebral trauma and may be an important contributory factor to a fatal outcome. Cerebral trauma *per se* affects the circulation to and from the cells, besides which shock, oedema, anaemia, hyperthermia and drugs all tend to one or other form of anoxia, further increasing the damage to the brain, the cells of which are extremely vulnerable to O_2 deprivation. Any of the different types of anoxia may affect the tissues of the brain. Thus anoxic anoxia results from congestion, oedema or consolidation in the lungs. Anaemic anoxia occurs where the head injury is accompanied by severe haemorrhage (as from scalp wounds) or in an atmosphere of carbon monoxide (as in mine accidents). Stagnant anoxia is present in conditions of shock but more particularly as a result of the congestion and stasis so frequently associated with severe head injuries. Histotoxic anoxia, in which there is a direct interference with tissue utilization of O_2 , could be particularly evoked by the undue administration of narcotics or anaesthetics. In the severely traumatized patient the combination of all the above factors plays no small part in the fatal outcome of the case. Depression of the blood O_2 can range from 5% to as much as 44% (Schnedorf *et al.*, 1940). The most permeable section of the small vessels appears to be at the point of transition between capillaries and venules. In stasis (possibly consequent to vasoparalysis—Scheinker, 1944) anoxic damage to the vessel wall increases its permeability, allowing the escape of the fluid and formed elements of the blood into the perivascular spaces, thus further compressing the vessels and causing a vicious circle of congestion and stasis. This is enhanced by the accumulation of abnormal metabolites in anoxic tissue causing the latter to absorb more fluid and accentuating the already existing oedema (Masserman, 1934).

(To be continued)

INFANTILE ECZEMA*

S. GORDON, M.B., Ch.B. (WITS.), M.R.C.P. (Ed.)

Johannesburg

The subject of infantile eczema should be approached with humility. There is so much about it that is not known and so much that passes as knowledge which is no more than the cumulative repetition of a hundred textbooks.

* This paper was read to a meeting of the Northern Transvaal Branch of the Medical Association of South Africa in Pretoria on 12 June 1951.

The word eczema has come down to us from antiquity and has unfortunately brought its antiquity with it, so much so that the whole subject is still encrusted with the barnacles of pseudo-medical terminology.

Dermatologists have laboured for many years to raise the status of eczema to the level of an ordinary medical phenomenon—often at cross purposes, often confusing

elaboration with clarification, and always a little aghast at their own temerity. All these labours have at times added a little to our knowledge, sometimes they only added to our understanding, and often they only served to bring our ignorance into sharper relief. On the basis of the little knowledge that we now have and with the aid of some legitimate inference, it is to-day possible to adopt, in outline, a classification of eczema to which future investigation will give substance and shape.

If we accept some such classification—and we must, if we are going to particularize about infantile eczema—we begin with a definition of eczema as an inflammatory reaction of the epidermis to any substance to which it has become sensitized. By the terms of this definition the choice of an eczema subject is not necessarily determined by his age, heredity, diathesis or constitution. Similarly the sensitizing substance is not exclusively an external contactant, or an ingested substance of known composition, or an endogenous toxin of unknown nature. All that is necessary for an eczema reaction to take place is that the eczema subject and the sensitizing substance should have met and developed a mutual antagonism. Henceforth, whenever they meet, they will recognize each other and display that antagonism in the form which has become familiar to us as an eczema reaction.

Based broadly on this definition is a classification of the eczemas into the following groups:—

1. *Exogenous Eczema.* Here the reaction depends on a known and demonstrable contactant.

2. *Infective Eczema.* In this the sensitizing substance is bacterial or fungal and produces the reaction also by external contact.

3. *Endogenous Eczema.* In this the sensitizer is either a known substance which has been ingested, injected or inhaled or, by inference, a circulating toxin of unknown nature and doubtful existence.

This classification meets with difficulties when it comes to the atopic eczemas, which are said to be more common in some families than in others and in which there is a multiple sensitivity which may be both endogenous and exogenous.

To a very large extent the clinical picture of an eczema shows a corresponding morphological division. Thus an exogenous chemical eczema is commonly diffuse and vesicular, an infective eczema is usually a sheet which increases by peripheral extension, and an endogenous eczema is discoid and papular; but this is no more than a rough differentiation and should not be regarded as a complete guide to etiology. The essential unity of the eczema reaction is best demonstrated by the sulphur drugs which can produce the same eczematous rash when taken by mouth or applied topically. It is also illustrated by the behaviour of certain external irritants, notably nail polish, which can produce a diffuse eruption in some individuals and discoid scaly patches in others.

Such a classification of eczema has at last hesitantly insinuated itself into the pages of conservative dermatology,¹ but the benefit of rational classification has not yet been accorded to infantile eczema. This is still heavily-handedly described as a single entity, nebulously attributed to endogenous toxins, allergens or atopins. This is oversimplification at its worst. To accept such a single etiology is to deny that an infant is subject to the same laws of antigen-antibody reaction as an adult. Percival,² who

goes a little further than other writers, admits the existence of other forms of eczema in infants and distinguishes between eczema in infants and infantile eczema. This is an unnecessary complexity. Once it is admitted that exogenous forms of eczema do occur in infants, then for all practical purposes every case of infantile eczema becomes an eczema in an infant and may belong to any of the groups that we recognize in adults.

Following the same sequence that we adopted for adult eczema, infantile eczema can now be discussed under the following headings:

Contact Eczema. As a baby does not use cosmetics and is not exposed to industrial hazards it escapes the two common sources of exogenous sensitization and the incidence of contact eczema in infants is almost negligible; almost, but not quite. F. A. Simon³ described a contact sensitivity to the mother's dandruff which could be demonstrated by patch tests. Gaul and Underwood⁴ described a generalized eczema reaction in a baby as a sequel to sensitization by footwear. A personal case is that of a baby sensitive to the mother's hand lotion. The importance of recognizing this group, however, lies in the fact that infants are daily exposed to the risk of sensitization by skillfully advertised ointments of questionable value and high sensitizing potential. As long as mothers will rub patent ointments into flea bites and pour drops into stuffy noses, the possibility of exogenous sensitization will always be with us. Moreover, the use of such ointments is liable to convert a mild bacterial or endogenous eczema into a severe chemical eczema.

Infective Eczema. It has been recognized for many years that some infantile eczemas start with a cradle cap, show an early tendency to affect the retro-auricular area, the axillae, and the groin, and appear as sharply marginated, bright red, oozing and crusted sheets. Misguided attempts have been made to bring these within the orbit of seborrhoeic eczema. As seborrhoeic eczema has only a mythical existence in the adult, and as infants do not suffer from greasy skins, or seborrhoea, this term can hardly be regarded as aptly descriptive. The recognition of this form of eczema as a bacterial infection and bacterial sensitization is a real advance in dermatology and the term seborrhoeic eczema, in general, should now be allowed to go to its final resting place and forgotten. Unfortunately, the victims of sulphosalicylic ointment therapy will have reason to remember it for some time to come.

There is a definite relationship between impetigo and eczema in infants. This is important because all forms of impetigo are common in infants. There is the acute impetigo contagiosa; there is the chronic streptococcal impetigo, which starts from a discharging ear; there is the fine scaly impetigo of the face and the more generalized scaly impetigo of the trunk which has been described as erythema streptogenes.⁵ Another form of impetigo, which appears as circumscribed, finely crusted, erythematous patches and is often mistaken for eczema was described by Adamson⁶ in 1908 and more recently by Crawford⁷ as chronic symmetrical impetigo. These forms of impetigo are mentioned here not only because they can be mistaken for eczema, but because in certain circumstances they become complicated by an eczema reaction. This brings us to the subject of auto-eczematization.

Under the term autosensitization Whitfield⁸ described

the appearance of widely distributed papulo-vesicular patches a number of days after an exacerbation of a chronic localized, lichenified eczema. Recently Cormia and Esplin⁹ described a similar phenomenon which they call auto-eczematization. Briefly their concept is that epidermal cells in a state of inflammation may become antigenic and produce widespread epidermal reactions. This process is enhanced by chemical irritation or mechanical trauma. Cormia and Esplin believe that liberation of antigens may be determined by bacterial infection and chemical sensitization of a limited area of skin. They suggest that clinically distinct entities, e.g. nummular eczema, atopic eczema and infective eczema may have this mechanism in common. In fact, they hope that this concept will unravel the enigma of eczema.

The application of this concept to infantile eczema is very obvious. Its most practical application is in this group of infective eczemas. It is only necessary to accept the hypothesis that epidermal cells altered by bacterial infection may acquire antigenic properties to have an explanation for the development of eczema as a sequel to an impetiginous infection of the skin. Add to this the dangers of auto-eczematization by chemical irritation and the trauma of scratching and one begins to wonder whether this sequence does not account for a large part of infantile eczema. Personally, I think it does.

Atopic Eczema. Under this heading may be conveniently grouped all eczema reactions, without discoverable exogenous causes. Here we come to the submerged part of medicine where a little speculation is permissible and is, indeed, inevitable. For the sake of simplicity atopy may be defined as an inherently low sensitivity threshold. By this definition persons belonging to certain families are liable to acquire a polyvalent sensitivity to a variety of sensitizers, or atopens, the effect of which is the development, among other sensitivity phenomena, of an eczema which appears early in life and may persist throughout childhood and into adult life. Clinically this form of eczema is a dry, papular, often lichenified eruption in somewhat ill-defined sheets affecting the face, the antecubital and popliteal areas, the ankles and the wrists. Etiologically this group is very ill-defined. It cannot even be regarded as purely endogenous, because sensitivity to contact with woollen garments is known to be a factor. However, to many dermatologists and pediatricians atopic eczema is almost synonymous with food allergy and calls for an immediate assault on the baby's diet. In terms of our definition of eczema it must be accepted that some eczema reactions are caused by food allergens, but this over-emphasis of food allergy would lose much of its background if the subject were reviewed and all the evidence based on skin allergy tests were finally and unhesitatingly discarded.

There are some perplexing questions in connection with food allergy to which the answer is not obvious. Firstly, if food allergy is a common cause of eczema in infants, why is it that urticaria, the more familiar manifestation of food allergy, is so uncommon in infants? Secondly, food allergy as a cause of eczema is now well recognized in adults. Why is it then that all the investigators in this field¹⁰ have found that the maximum incidence is as an eruption of the hands and not of the areas commonly

affected in atopic eczema? Thirdly, what is that phenomenon which takes place at the age of two or three and which produces a sudden and permanent disappearance of the sensitivity in the majority of infantile eczemas? In the present state of our knowledge there is not much more that can be said about atopic eczema, except to refer again to an observation that has been made before, that atopic eczema has many points of similarity with dermatitis herpetiformis.¹¹ Dermatitis herpetiformis was described by Duhring in the last century¹² and since then it has gradually come to mean an itchy, pigmented, bullous eruption, but this imposes a limit on Duhring's work which Duhring never intended it to have. Duhring also described other types of dermatitis herpetiformis, including a papular eczematous type. Between this papular type and the papular type of eczema the dividing line becomes so blurred that it is a reasonable speculation that dermatitis herpetiformis, atopic eczema and auto-eczematization are either the same or closely related phenomena. The importance of this will be seen when we come to treatment, but before we do that it might be convenient at this stage to mention two other conditions which hover on the horizon to catch the unwary. One is prickly heat. This is an itchy eruption of fine translucent papules on the covered parts of the body. The affected skin feels as if it had been dusted with fine sand. It is dealt with very simply by stopping the baby's bath for a week or so and using a mild salicylic lotion. The other is a non-itchy fine papular eruption which often follows an upper respiratory infection. It is regarded as a streptococcal and responds to sulphonamide therapy.

The treatment of infantile eczema follows on naturally from the classification which has been adopted. In the contact-type exogenous eczema, all that is necessary is to eliminate the offending allergen and apply a bland lotion. It must be emphasized that general sensitization is always imminent and that the thoughtless application of 'panacea' ointments to this type of eczema almost amounts to malpraxis. The infective type of eczema is dealt with primarily by the topical application of antiseptics of low sensitizing potential. The applications that deserve special mention are Tyrothricin solution 0.1% which is used as a wet dressing, and Vioform in 3% concentration as a lotion, cream or paste, depending on the stage of the eczema reaction. While not a great adherent of polychromatic dermatology, I must admit that some of the dyes, e.g. 1% brilliant green, have their uses in this group. After a period of antiseptic treatment, when the eruption has largely subsided and the skin has become dry and scaly, it is often useful to apply a crude coal tar paste. Incidentally, applications of crude coal tar paste have a not undeserved reputation in all forms of eczema, when this particular stage has been reached.

In atopic eczema the external treatment is only of secondary importance. As in the infective type, the application of some antiseptic in the presence of secondary impetigo is helpful and the application of crude coal tar may relieve the itching, but this does not influence the atopic state.

A new approach to the treatment of the atopic state has emerged as a result of the observation of the similarity between atopic eczema and dermatitis herpetiformis. The

etiology of dermatitis herpetiformis is as much a mystery as the etiology of eczema. What is known about dermatitis herpetiformis is that the eruption can be kept suppressed indefinitely by arsenic or Sulphapyridine, or para-aminobenzoic acid. It is also known that suppression of the condition for long periods will in many cases bring about an eventual cure.^{1,2}

After an extensive trial in cases of atopic eczema and auto-eczematization it can now be added that the similarity between dermatitis herpetiformis and atopic eczema is even more striking in the field of therapeutics. Without trying to explain the rationale of this treatment, without claiming infallibility for this method, it can now be stated that the administration of Sulphapyridine or Pabavel has proved of inestimable value in the management of atopic eczema in infancy. Until more is known about the anti-eczematous properties of these substances, it is only possible to say that their action on the epidermis is comparable with the action of antihistaminic drugs on the dermis.

IN MEMORIAM

DR. J. S. MORTON

Dr. James Stevenson Morton, the doyen of the medical profession on the East Rand, died on Saturday, 21 July 1951, at the age of 84.

He was born in Northern Ireland and qualified at Trinity College, Dublin, obtaining the M.B. Ch.B. and subsequently proceeded M.D. He came to South Africa in 1894 as Medical Officer to the Fattar Group of Mines, a position he held jointly with the late Dr. J. McNeillie. During the Boer War he took part in the siege of Kimberley. At the end of this war he returned to Boksburg, becoming Medical Officer to the E.R.P.M. Ltd. He was also employed as Medical Officer to Brakpan Mines from 1906 to 1908 and the Geldenhuys Deep for a short period.

For many years he held the post of Senior Honorary Physician to the Boksburg Hospital and subsequently to the Boksburg-Benoni Hospital, retiring from this position in 1936. He was the Medical Officer of Health to the Boksburg Municipality for a number of years. For many years he was a valued member of the Boksburg-Benoni Hospital Board.

He was a keen member of the B.M.A. and numbered amongst his friends many of the leading surgeons of Johannesburg. In 1944 he was made an Honorary Member of the B.M.A. after 50 years of unbroken membership and subsequently he was similarly honoured by the local East Rand Division and East Rand Branch of the M.A.S.A. On the latter occasion he delivered a stimulating address on 50 years of practice on the East Rand, showing a keen appreciation of modern developments and trends in medicine.

He held high office in the Masonic world; was a keen and able golfer and his company was much sought after on the golf course. In his later years he became a keen bowler and proved himself adept at this game.

- ### REFERENCES
1. Mitchell Heggis, G. B. (1950): *Modern Practice in Dermatology*, p. 188. London: Butterworth & Co.
 2. Waiker, N. and Percival, G. H. (1947): *An Introduction to Dermatology*, pp. 91, 117. Edinburgh: E. & S. Livingstone Ltd.
 3. Simon, F. A. (1945): *Arch. Dermatol. Syph.*, **51**, 402.
 4. Gaul, L. E. and Underwood, G. B. (1949): *Arch. Dermatol. Syph.*, **60**, 649.
 5. Dobes, W. L. and Jones, J. (1946): *Arch. Dermatol. Syph.*, **53**, 107.
 6. Adamson, H. G. (1908): *Brit. J. Dermatol.*, **20**, 109.
 7. Crawford, S. (1949): *Arch. Dermatol. Syph.*, **59**, 55.
 8. Whitfield, A. (1930): *Trans. 8th Internat. Dermatol. Cong.*, Copenhagen.
 9. Cormia, F. E. and Esplin, B. M. (1950): *Arch. Dermatol. Syph.*, **61**, 931.
 10. Rowe, A. H. (1946): *Arch. Dermatol. Syph.*, **54**, 683.
 11. Flood, J. M. and Perry, D. J. (1946): *J. Invest. Dermatol.*, **7**, 309.
 12. Gordon, S. and Loewenthal, L. J. A. (1949): *Brit. J. Dermatol.*, **61**, 359.
 13. Duhring (1893): *Selected Monographs on Dermatology*. London: New Sydenham Society, **143**, 184.
 14. Peterkin, G. A. G. (1951): *Brit. J. Dermatol.*, **1**, 63.

He continued in practice up to his last illness. A man of kindly disposition, he was esteemed by patients and colleagues alike.

G. B. M.

A correspondent writes: The death occurred in Boksburg, Transvaal, on 21 July 1951, of James Stevenson Morton, M.D., D.P.H., aged 84 years. He was the elder son of Mr. James Morton, of Mount Pleasant, Gilford, Northern Ireland, and was born on 18 April 1867. He was educated at the Belfast Academy and then at Trinity College, Dublin, where he obtained his Medical degree. He proceeded to 'The Rotunda' for midwifery, and later took his D.P.H. at Cambridge.

Dr. Morton came out to South Africa in 1895, arriving in Boksburg on 24 December of that year. He was granted a licence to practise Medicine in the Transvaal Republic under President Kruger. He joined the British Forces in 1899 at the outbreak of the Boer War, and was in Ladysmith throughout the historic siege. After the Relief, he was appointed to a Hospital Ship plying between South Africa and Britain where he served until the Armistice in 1902, when he returned to Boksburg.

He was appointed M.O.H. for Boksburg in 1905 and served in that capacity until 1936. He did honorary work for the Boksburg-Benoni Hospital from the time of its inception until very recently, and apart from his general practice, was one of the E.R.P.M. (East-Rand Proprietary Mines) doctors.

Dr. Morton was a prominent member of the Masonic Lodge in Boksburg and took a keen interest in all public affairs. He continued in general practice until his last illness, in June this year, having been practising in Boksburg for 56 years. He leaves a widow and one daughter, his only son having been killed in action in the Royal Navy in 1945.

PASSING EVENTS: IN DIE VERBYGAAN

We regret to record the death of Dr. M. A. Diemont, of Stellenbosch, who passed away on 1 September 1951.

We regret to record the death of Dr. J. A. Weir, a general practitioner of Rondebosch, Cape, who passed away on 4 September 1951.

Dr. Percy Helman of Cape Town has been successful in the Primary Examination for the Fellowship of the Royal College of Surgeons, England.

TENTH INTERNATIONAL CONGRESS OF DERMATOLOGY: 1952

The Academic Committee is now prepared to receive applications from dermatologists who wish to read short papers at

this Congress. Speakers will have 15 minutes in which to present their papers. In the majority of instances this will be followed by a discussion not exceeding 15 minutes.

Chairmen of Meetings will be asked to insist on the 15-minute schedule.

Titles of communications and an accurate note concerning the name, appointments and degrees of the applicant should be forwarded to the Congress through Dr. J. Frootko, 1006 Medical Centre, Jeppe Street, Johannesburg, by 1 January 1952.

Dr. C. C. Pretorius is terug van sy nagraadse studies oorsce, en spesialiseer nou as verlos- en vrouesiektekundige te Transvaliegebou 416, Sentraalstraat, Pretoria.



CORTOGEN ACETATE OPHTHALMIC SUSPENSION

IN A SPECIALLY STABILIZED AQUEOUS VEHICLE

CORTISONE ACETATE SPECIFICALLY FOR THE EYE

CORTOGEN Acetate Ophthalmic Suspension gives safe, simple and often dramatically effective control of inflammatory and allergic reactions in the anterior segment of the eye. Its use is virtually free of irritation to the tissues of the eye. CORTOGEN Acetate Ophthalmic Suspension is indicated in allergic conjunctivitis, blepharitis, phlyctenular conjunctivitis, keratitis, iritis, and many other disorders of the eye.

CORTOGEN Acetate Ophthalmic Suspension in dropper bottles of 3.75 cc. 5 mg. Cortisone Acetate per cc.

Schering

CORPORATION • BLOOMFIELD, N.J.

Sole Distributors • SCHERAG (PTY.) LTD., P.O. BOX 7539 • JOHANNESBURG



A Calamine-type Lotion containing 'Benadryl'

CALADRYL

'Caladryl' is a smooth, creamy lotion containing 1 per cent of the anti-histamine and anti-pruritic agent 'Benadryl' with calamine, camphor and glycerin. The lotion has emollient properties but is non-greasy.

'Caladryl' is effective in allaying the burning and irritation of sunburn and in relieving itching due to insect-bites. It also relieves pruritus associated with urticaria, dermatitis, minor skin affections, measles and chicken-pox, and is indicated in all forms of cosmetic allergy.


For infants 'Caladryl' may be used for napkin-rash and teething-rash.

In 4 fluid ounce bottles



**PARKE, DAVIS
& COMPANY, LIMITED**
HOUSLOW, near LONDON

Further information from any branch of LENNON LTD.



Ready-to-inject

Crysticillin Suspension

SQUIBB Procaine Penicillin G in Aqueous Suspension

**Stable for 1 year at room temperature
no refrigeration required**

SUPPLIED IN 5- AND 10-DOSE VIALS
containing 300,000 units per c.c.

*Squibb Crysticillin is also available
in various dry forms as follows:*

- 1 c.c. Fortified Crysticillin
(400,000 units per c.c.)
- 5 c.c. Crysticillin Dry 1½ million
units (300,000 units per c.c.)
- 5 c.c. Fortified Crysticillin 2
million units (400,000 units per c.c.)
- 10 c.c. Crysticillin Dry 3 million
units (300,000 units per c.c.)

"Crysticillin" is the
registered trade mark
of E. R. Squibb & Sons.

SQUIBB

A LEADER IN PENICILLIN RESEARCH AND MANUFACTURE

PROTEA PHARMACEUTICALS LIMITED

● Further information and literature available from

P.O. Box 7793 Johannesburg

Phone: 33-2211

GRACEFUL women, distinguished men,
soft lights, scintillating conversation—
such is the atmosphere in which Viceroy
Cigarettes are consistently chosen.

Wills's
VICEROY



PLAIN.

CORK & FILTER



clinical efficiency

The germicidal efficiency of 'Dettol' remains high even in the presence of blood, pus and wound debris. This property, coupled with its wide margin of safety, makes 'Dettol' invaluable for use in emergencies, not only by you, but in the less qualified hands of others who in emergency might have to render first aid.

DETTOL

THE MODERN ANTISEPTIC

RECKITT & COLMAN (AFRICA) LTD., P.O. BOX 1097, CAPE TOWN

34

1805-2E

VALUABLE BOOK FREE!

ARE YOU PREPARING FOR ANY MEDICAL,
SURGICAL, or DENTAL EXAMINATION?

Send Coupon below for our valuable publication

"Guide to Medical Examinations"

PRINCIPAL CONTENTS:

The Examinations of the Conjoint Board.
The M.B. and M.D. Degrees of all British Universities.
How to pass the F.R.C.S. Exam.
The M.S. Lond. and other Higher Surgical Examinations.
The M.R.C.P. London.
The D.P.H. and how to obtain it.
The Diploma in Anaesthetics.
The Diploma in Psychological Medicine.
The Diploma in Ophthalmology.
The Diploma in Laryngology.
Diploma in Radiology.
The D.R.C.O.G. and M.R.C.O.G.
The Diploma in Child Health.
Do not fail to get a copy of this Book before commencing preparation for any Examination. It contains a large amount of valuable information. Dental Exams. in special Dental Guide.

SEND FOR YOUR COPY NOW!

The Secretary,

MEDICAL CORRESPONDENCE COLLEGE

19 Welbeck Street, Cavendish Square, London W.1.

SIR,—Please send me a copy of your "Guide to Medical Examinations" by return.

Name

Address
Examination in
which interested

S.A.M.J. South African Offices: P.O. Box 2239, Durban Natal

Showell's



Suture Needles

Sole Distributors for the Union of South Africa

Chas. F. Thackray

301-303 Boston House, Strand St. (P.O. Box 816) CAPE TOWN

23 Orion House, 235 Bree St. (P.O. Box 2726) JOHANNESBURG

Provincial Administration of the Cape of Good Hope

HOSPITALS DEPARTMENT

HONORARY APPOINTMENT

Applications are invited from registered medical practitioners for the following post:

Honorary Surgical Registrar at the Victoria Hospital, Wynberg.

The appointment will be for the remainder of 1951.

Applications containing particulars of age, qualifications, experience, etc., with copies of recent testimonials should be forwarded to the undersigned by Noon, 29 September 1951.

L. Welham

Branch Representative

Hospitals Department
Industry Building
58 Loop Street
Cape Town

4734

Transvaalse Provinsiale Administrasie

VAKATURES BY PUBLIEKE HOSPITALE

Aansoek word ingewag van kandidate met geskikte kwalifikasies vir die onderstaande pos by die Ontdekkersgedenks-hospitaal, P.K. Florida in die Transvaal.

Aansoek met gerig word aan die Verantwoordelike Geneesheer van die Ontdekkersgedenks-hospitaal en moet volle besonderhede bevat aangaande die ouderdom, professionele, akademiese en taalkwalifikasies, ondervinding en huweliksstaar van die applikant en moet voorts 'n aanduiding bevat van die vroegste datum waarop diens aanvaar kan word.

(Ontdekkersgedenk - Deeltydse Algemene Praktisyn Chirurg, £225 p.a.) moet hospitaal daglikse besoek.

Van die persone wat aangestel word, sal ver wag word om bevredigende sertifikate in te dien.

Aansoekvorms is verkrygbaar van die Provinsiale Sekretaris, Departement van Hospitaaldienste, Posbus 383, Pretoria.

Die sluitingsdatum van aansoek vir die pos is 2 Oktober 1951. (30923)

Royal College of Surgeons of England

PROFESSORSHIP OF HUMAN AND COMPARATIVE ANATOMY

The Council invites applications for the Sir William Collins Professorship of Human and Comparative Anatomy.

The salary will be from £2,000 to £2,500 per annum according to age and academic status, with arrangements for membership of the Federated Superannuation System for Universities.

The election is an open one, but the Council reserves the right to invite individual applications.

Particulars and terms of appointment may be obtained from the Secretary. Applications with the names of two referees must reach the College not later than 30 November 1951.

Kennedy Cassels

Lincoln's Inn Fields
London W.C. 2, England

Secretary
4 RC/C 58

Vakature vir Mediese Beemple

Die aadig word gevestig op 'n advertensie in die *Staatskoerant* van 14 September 1951, waarby aansoek om aanstelling gevra word, op kontrak, in 'n vakante pos van Mediese Beemple in die personeel van die Gesondheidsinstituut vir Gesin en Smelewing, Clarwood, Durban.

Die salarisskaal aan die pos verbonde, is £720 + 30 - 900 + 40 - £1,020, plus 'n lewenskostetoelae van £256 per jaar in die geval van getroude en £80 per jaar in die geval van ongegetroude amptenare.

Aansoekvorms is verkrygbaar van die Sekretaris van Gesondheid, Posbus 386, Pretoria. (30872)

The Medical Association of South Africa Die Mediese Vereniging van Suid-Afrika

AGENCY DEPARTMENT: AGENTSAP AFDELING

KAAPSTAD: CAPE TOWN

Posbus 643, Telefoon 2-6177: P.O. Box 643, Telephone 2-6177

PRAKTYKE TE KOOP: PRACTICES FOR SALE

(644) Durban Central. Mainly Indian and Native cash practice. Average annual gross income £1,235. Premium of £500 required for goodwill, inclusive of furniture and fittings and drugs. Terms may be arranged.

(805) Transkei Native and D.S. practice. Near large town. House and surgery for sale, purchaser preferably bilingual.

ASSISTENTE PLAASVERVANGERS VERLANG ASSISTANTS LOCUMS REQUIRED

(754) For Transkei Native and D.S. practice. Initial salary £60 p.m. and all found. Definite view to partnership after trial period. Single man preferred.

(724) Cape Town. Excellent opportunity for assistant in Northern suburbs. Salary, etc., to be arranged.

(744) Noordwest. Assistent vir praktyk met klein private verpleeginrigting - £65 p.m. plus vry losies. Kar beskikbaar.

(792) Cape Town Southern suburb. Middle- and better-class European practice. From 1 December for 4½ weeks. £2 2s p.d. plus 10s. car allowance p.d. plus board and lodging. Must have done some general practice previously. Very little night work.

(723) Assistantship in hospital town near Cape Town. Salary £75 p.m. and car provided. Dwelling available. Possibility of permanent appointment.

(798) Cape Town. From 14 October for three weeks. £3 3s. per day. Locum to use his own car and live at principal's home.

(801) South-Eastern Cape. Large partnership in hospital town requires a locum as from 1 October. Preferably Afrikaans and married. £2 2s. per day plus all found, allowance for use of car. Possibility of long-term appointment.

JOHANNESBURG

Medical House, 5 Esselen Street. Telephone 44-9134-5, 44-0817

Mediese Huis, Esselenstraat 5. Telefoon 44-9134-5, 44-0817

PRAKTYKE TE KOOP: PRACTICES FOR SALE

(Pr S30) Johannesburg Partnership practice plus Solus practice. Mainly non-European. Present income £3,600 p.a. Premium for quick sale £1,250.

(Pr S31) O.V.S.-praktyk. Goeie geleentheid vir algemene geneesheer met aanleg vir snywerk. Alle fasiliteite. Medisyne word aangemaak. Moet tweetaalig wees. Jaarlikse inkomste £2,400. Eienaar gaan verder studeer. Premie vir klandise-waarde, instrumente en voorrade, £1,500. Een maand introduk-sie sal gegee word.

ASSISTENTSKAP VERLANG: ASSISTANTSHIP REQUIRED

(A W52) Assistantship in established practice in S. Rhodesia required by married South African doctor. Qualified two years.

ASSISTENT/PLAASVERVANGER VERLANG ASSISTANT/LOCUM REQUIRED

(L V112) Johannesburg from 18 December 1951 to 18 January 1952. £2 12s. 6d. p.d. plus all found and petrol and oil paid.

MEDICAL EQUIPMENT

(I 017) Leitz microscope £45

(I 018) What offers? Complete set 'British Encyclopaedia of Medical Practice' plus all editions of 'Medical Progress' to 1950. Condition as new.


(I 019) Zeiss microscope. Condition as new. £55.

(I 020) "Standby" model Baumanometer. £10.

(I 021) Portable Baumanometer. £3.

(I 022) Klinostik Auroscope with Ophthalmoscopic attachment. £3.

(I 023) Heavy based Irrigator stand, height adjustable, complete with glass flask and hook to carry vacolitre flasks. £7.



PARENTERAL administration of Luminal promptly tones down nervous system excitement. Whether manifested as convulsions, psychic agitation or pernicious vomiting, nervous overactivity is controlled profoundly and for prolonged periods with Luminal in adult doses of from $\frac{1}{4}$ to 3 grains.

By prescribing Luminal, certainty of therapeutic result and accurate dosage is assured. For, when Luminal is specified in your prescription there is no valid excuse for the substitution of something else. Hundreds of published clinical reports testify to its therapeutic efficiency. Your preference for the pioneer brand of phenobarbitone is therefore well-founded.

LUMINAL

The pioneer brand of phenobarbitone

BACKED BY MORE THAN 30 YEARS OF EXPERIENCE



Winthrop Products
(PTY.) LTD.

Box 4186
CAPE TOWN

Box 9536
JOHANNESBURG

Box 2461
DURBAN

Desoxyn

HYDROCHLORIDE

*Methamphetamine
Hydrochloride, Abbott
tablets,
elixir and
ampoules*

Indications

Desoxyn Hydrochloride is indicated for oral administration in the treatment of narcolepsy, in cases of mild depression and for temporary use as a mental stimulant, where valid indications are present.

Favourable results have also been reported following the use of d-desoxy-ephedrine hydrochloride as an adjunct in the treatment of postencephalitic Parkinson's syndrome, chronic alcoholism, cerebral arteriosclerosis and generally in conditions for which amphetamine sulfate has been of benefit.

LONGER EFFECT

Average duration from an oral dose of 10 mg. is 6 to 12 hours, with up to 26 hours reported. Following parenteral administration to maintain blood pressure during and after surgery, satisfactory levels usually are sustained several hours.

FEWER SIDE-EFFECTS

Recommended dosages rarely cause undesirable reactions. If side-effects do occur, the dose may be reduced, or the drug withdrawn and Numbatal given as a sedative. While it is not habit-forming, it of course should not be used over prolonged periods as a substitute for sleep and rest.

QUICKER ACTION

Rapid absorption provides stimulation in 20 to 30 minutes following oral dosage. Intravenously, DESOXYN produces a rise in systolic pressure almost at once, maximum effect in 2 to 3 minutes. Intramuscular injection results in a rise in 2 to 10 minutes, with maximum effect in 4 to 20.

SMALLER DOSAGE

Greater potency by weight means that a smaller dose may be used—from 2.5 to 10 mg. daily for oral administration is usually sufficient. In emergencies, 10 to 15 mg. may be given intravenously; for non-emergency use, 15 to 30 mg. intramuscularly, depending on condition of the patient.

How supplied

Desoxyn Hydrochloride Tablets, 2.5 mg. are supplied in bottles of 25 and 100 (List No. 3488).

Desoxyn Hydrochloride Elixir, containing 20 mg. per fluidounce (2.5 mg. per fluidrachm), is supplied in bottles of 1 pint (List No. 3703).

Desoxyn Hydrochloride Ampoules, 20 mg. per c.c. are supplied in 1-c.c. size, in boxes of 6 (List No. 3378).



ABBOTT LABORATORIES, S.A. (Pty.) Ltd.

JOHANNESBURG · CAPE TOWN · DURBAN